



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

January 16, 2018

Ye Liu  
Regulatory Specialist  
Stepan Company  
22 Frontage Road  
Northfield, IL 60093

Subject: Label Amendment – Revisions to precautionary statements, alignment with CSF and other minor modifications  
Product Name: SC-5:256N  
EPA Registration Number: 1839-235  
Application Date: November 2, 2017  
Decision Number: 535813

Dear Ms. Liu:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2  
EPA Reg. No. 1839-235  
Decision No. 535813

with FIFRA section 6. If you have any questions, please contact Tara Flint via email at [flint.tara@epa.gov](mailto:flint.tara@epa.gov) or Eric Miederhoff at [Miederhoff.eric@epa.gov](mailto:Miederhoff.eric@epa.gov).

Sincerely,

A handwritten signature in blue ink that reads "E. Miederhoff". The signature is written in a cursive style with a large, stylized initial "E".

Eric Miederhoff  
Product Manager 31  
Regulatory Management Branch I  
Antimicrobials Division (7510P)  
Office of Pesticide Programs

Enclosure

**ACCEPTED**

01/16/2018

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 1839-235

# SC-5:256N

## FRONT PANEL

### ACTIVE INGREDIENTS

Didecyl dimethyl ammonium chloride ..... 19.596%  
Alkyl (50% C<sub>14</sub>, 40% C<sub>12</sub>, 10% C<sub>16</sub>)..... 13.064%  
dimethyl benzyl ammonium chloride

**OTHER INGREDIENTS** .....67.34%  
**Total**.....100.000  
%

**NET CONTENTS(:)** XXX(oz)(ounces)(gallon(s))(gal(s))

**KEEP OUT OF REACH OF CHILDREN**

# DANGER

FIRST AID	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
<b>If in Eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on Skin or Clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call poison control center or doctor for treatment advice.</li> </ul>
<b>If Inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If Swallowed:</b>	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything to an unconscious person.</li> </ul>
<b>(HOTLINE NUMBER)</b> You may contact (xxx) xxx-xxxx for emergency medical treatment information.)	

### NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

See (side) (back) panel for additional precautionary statements.

## FRONT, BACK OR SIDE PANEL

EPA REG. NO. 1839-235  
EPA EST. NO. 1839-IL-01 (XXX-XX-XX)

### STEPAN COMPANY

22 West Frontage Road  
Northfield, IL 60093

Page 1 of 20  
Amended 11/02/2017  
EPA Reg. No 1839-235

Bracketed information is denoted as: {insert as noted }, ( optional text ), [ required qualifier ] , <note>.  
The term "this product" used throughout this document may be replaced with the marketed product brand name.  
The term "x" is a place holder for numerical digits.  
Punctuation and plural/singular word forms may be adjusted to allow for grammatical correctness.  
Appropriate metric or imperial unit may be added as optional, supplemental information.

## **PACKAGING CLAIMS**

- Bonus (pack)(size)
- (Extra) value (pack)(size)
- (Dual)(X) pack
- Twin pack

## **MARKETING CLAIMS**

### **General**

- Clean(er)(ing)
- Disinfect(ant)(ing)
- Sanitize(r)
- Detergent
- Virucide\*
- Fungicide (against pathogenicfungi)
- Mildewcide
- Mildewstat
- Deodorize(r)(s)
- Kills Germs
- Kills Mold & Mildew
- pH Neutral
- Effective cleaning (strength)(power)
- Concentrate(d)
- Phosphorous free
- Free of phosphates

- |  |
|--|
| <ul style="list-style-type: none"><li>• Citrus canker control (<i>Not for Use in California</i>)</li></ul> |
|--|

- This product is for use (in)(on) {insert use site or surface from page 5-7}.
- This product is a phosphate-free, pH neutral formulation designed to provide effective cleaning, deodorizing, and disinfection for {insert use site or surface from page 5-7} (where housekeeping is of prime importance in controlling cross contamination).
- This product saves time, labor, and cost.
- This product is specially formulated to be non-dulling to floor finishes and requires no rinsing.
- Effective in the presence of 5% serum contamination
- (For) indoor/outdoor (use)
- Contains no phosphorous nor nonyl phenol ethoxylate detergents
- Effective in 5 minutes in presence of Hard Water and 5% serum
- This product is formulated to a neutral pH, yet provides effective cleaning strength that will not dull high gloss floor finishes.
- (This product is) an economical concentrate (that can be used with a mop and bucket, trigger sprayers or sponge)

## **DISINFECTION CLAIMS**

- This product or product name) is a one step neutral disinfectant that is effective against a broad spectrum of bacteria.
- Formulated for Effective Farm Premise Disinfection
- Formulated for Effective Poultry Premise Disinfection
- Formulated for Effective Swine Premise Disinfection

- Formulated for Effective Veterinary Practice/Animal Care/Animal Laboratory Disinfection
- One-Step Disinfectant, Germicidal Cleaner and Deodorant
- (5)(Five) minute hospital disinfectant
- Kills 99.9% of bacteria in (just) (5)(five) minutes\*
- This product is a 5 minute hospital disinfectant (effective in the presence of hard water and organic serum)
- Kills Pandemic 2009 H1N1 Influenza A virus (formerly called swineflu)
- Kills {insert pathogen from page 7-8}
- Disinfects against {insert pathogen from page 7-8}
- For large areas such as {insert use site or surface from page 5-7 } this product is designed to (clean and disinfect)(provide both general cleaning and disinfecting)
- (This product is) a no rinse neutral disinfectant cleaner that disinfects, cleans and deodorizes non-food contact surfaces in one (easy)(labor saving) step.
- Hospital disinfectant
- EFFICACY TESTS HAVE DEMONSTRATED THAT this product IS AN EFFECTIVE BACTERICIDE\*, FUNGICIDE, VIRUCIDE\*\*\* AND MILDEWCIDE IN WATER UP TO 200 PPM HARDNESS (AS CaCO<sub>3</sub>) IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).

#### *Optional Influenza Claims*

- Respiratory illnesses attributable to Pandemic 2009 H1N1 are caused by Influenza A virus. This product (Product Name) is a broad-spectrum hard surface disinfectant that has been shown to be effective against (Influenza A virus tested and listed on the label) and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).
- This product (Product Name) has demonstrated effectiveness against Influenza A virus and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 Influenza A virus.
- This product (Product Name) has demonstrated effectiveness against (Influenza A virus tested and listed on the label) and is expected to inactivate all Influenza A viruses including Pandemic 2009 H1N1 (formerly called swine flu).
- Kills Pandemic 2009 H1N1 Influenza A virus (formerly called swineflu).
- Kills Pandemic 2009 H1N1 Influenza A virus.

### **SANITIZATION CLAIMS<sup>2</sup>**

- Kills 99.9% of bacteria in (just) 15 seconds\*\*
- Mushroom farm sanitization [*Not for Use in California*]
- (Acts as) an effective sanitizer
- Formulated for sanitization
- Sanitizing (formula)(formulation)
- 

### **FUNGI / MOLD / MILDEW CLAIMS**

- (Inhibits)(prevents)(stops) the growth of mold and mildew
- Leaves {insert use site or surface from page 5-7 } (smelling) clean and fresh
- (This product ) will not scratch {insert use site or surface from page 5-7 }
- Kills 99.9% of (fungus)(fungi)(mildew)(mold) in (just) 10 minutes<sup>#</sup>
- Kills 99.9% of (fungus)(fungi) in (just) 5 minutes<sup>#</sup>
- Kills 99.9% of (fungus)(fungi)(mildew)(mold)<sup>#</sup>
- Kills (fungus)(fungi)(mildew)(mold)
- Kills athlete's foot fungus

- Kills fungus causing (ringworm)(ringworm of the foot)#
- Kills Trichophyton mentagrophytes (, the cause of (ringworm)(ringworm of thefoot))

## **DEODORIZING & CLEANING CLAIMS**

- Deodorizing Formula
- This product inhibits the growth of mold and mildew and their odors ((in)(on) {insert use site or surface from page 5-7 }) when use asdirected
- This product is a broad-spectrum germicidal cleaner and deodorant that is effective at 1:256 (0.5 ounces per gallon) dilution in the presence of 200 ppm hard water (calculated as CaCO<sub>3</sub>) plus 5% organic blood serum.
- This product deodorizes areas that are hard to keep (smelling) fresh, such as {insert use site or surface from page 5-7})
- This product deodorizes {insert use site or surface from page 5-7 } (,and other areas which are prone to odors caused bymicroorganisms)
- (When used as directed) this product (deodorizes)(will deodorize) surfaces in toilet areas, garbage cans and garbage storage areas, and other places where bacterial growth can cause malodors
- (Effective)(powerful)(quick)(superior) cleaningstrength
- Can be used with a (mop and bucket)(triggersprayer)(or)(sponge)

## **CLAIM QUALIFIERS**

\* This product is an effective bactericide against Staphylococcus aureus (ATCC 6538), Pseudomonas aeruginosa (ATCC 15442), (and) Salmonella enterica (ATCC 10708), {and insert bacteria from the disinfection section on page 7-8} on hard, non-porous, non-food contact surfaces in 5 minutes.

\*\* This product (sanitizes in 15 seconds)(is an effective sanitizer) against Staphylococcus aureus (ATCC 6538), {and insert bacteria from the sanitization section on page 7-8} on hard, non-porous, non-food contact surfaces.

\*\*\* This product (is an effective virucidal in 5 minutes)(is an effective virucide) against [insert viruses from page 7-8] on hard, non-porous, non-food contact surfaces in 5 minutes.

# This product is an effective fungicide against Trichophyton mentagrophytes and mildewcide against Aspergillus niger in 10 minutes, and Candida albicans in 5 minutes on hard, non-porous, non-food contact surfaces.

<sup>1</sup> Rinse all equipment that comes in prolonged contact with skin before reuse with clean warm water (about 120°F), and allow to air dry. (Caution: Cleaning at the 120°F temperature will avoid overheating and distortion of the personal safetyequipment that would necessitate replacement.)

<sup>2</sup> Non-food contact surfaces and use site(s) only

<sup>3</sup> [This product or product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body.] or [This product or product is not for use on critical and semi-critical medical device surfaces.]

## **SURFACES & USE SITES**

Product label may list surface or site and/or a graphic depicting the hard, non-porous surfaces below.  
 <Note: When used as a sanitizer, this product is for use on non-food contact surfaces only.>

### **Hard, non-porous surfaces:**

Appliance Exteriors	Glazed Porcelain	Restroom Fixtures
(Bath)tubs	Goggles <sup>1</sup>	Salad Bar Sneeze Guards
Bed Frames	Half Mask Respirators <sup>1</sup>	Sealed Granite
Bidets	Hard hats <sup>1</sup>	Sealed Marble
Bins	Inflatable Advertising Structures	Shelves
Cabinets	Inflatable Advertising Structures	Shower (doors)(curtains)
Cages	Inflatable play structures	Shower Stalls
Cages	Kennels	Sinks (Basins)
Carts	Laminate	Sports Equipment
Chairs	Light Switches	Stainless Steel <sup>3</sup>
Chrome	Litter Boxes	Stovetops
Coolers	Metal <sup>3</sup>	Tables
Desks	Metal Blinds	Telephones
Diaper Pails	Microwave (oven) exteriors	
Doorknobs (and pulls)	Mirrors	Toilets
Counters (Countertops)	Non-wood Furniture	Tools
Drinking Fountains	Obstacle Course Play Equipment	Urinals
Ear Muffs <sup>1</sup>	Obstacle Course Play Equipment	Vanity Tops
Exercise Equipment	Outdoor Patio Furniture	Vinyl
Exercise Mats	Patient Chairs <sup>3</sup>	(Washable) Walls
Face Shields <sup>1</sup>	Personal Protective Safety Equipment <sup>1</sup>	Wheelchairs <sup>3</sup>
Faucets	Plastic	Whirlpool Interiors
Floors	Plastic Laundry Baskets	Windows
Freezer Exteriors	Plastic Mattress Covers	Wrestling Mats
Full Face Breathing Apparatus <sup>1</sup>	Protective Headgear <sup>1</sup>	(Barber)(Salon) tools
Garbage(cans)(pails)	Racks	Scissors
Gas Masks <sup>1</sup>	Recycling Bins	Manicure/pedicure instruments
Glass	Refrigerator Exteriors	
Glazed Ceramic Tile	Remote Controls	

### **Use Sites:**

Airplanes	Food Processing Plants <sup>2</sup>	Operating Rooms <sup>3</sup>
Ampitheaters	Food Service Establishments <sup>2</sup>	Pet Animal Quarters
Animal Life Science Laboratories	Funeral Homes	Pet Shops
Athletic Facilities	Health Clubs	Play Grounds

Bracketed information is denoted as: {insert as noted }, ( optional text ), [ required qualifier ] , <note>.  
 The term "this product" used throughout this document may be replaced with the marketed product brand name.  
 The term "x" is a place holder for numerical digits.  
 Punctuation and plural/singular word forms may be adjusted to allow for grammatical correctness.  
 Appropriate metric or imperial unit may be added as optional, supplemental information.

Automotive Garages	Hog Farms	Poultry Farms
Bars <sup>2</sup>	Hospitals <sup>3</sup>	Public Eating Establishments <sup>2</sup>
(Barber)(Beauty)shops	Hotels	Recreation(al) Facilities
Bathrooms	Household Bathrooms	Restaurants
Boats	Household Kitchens	Restrooms
(Breeding)(Grooming) Establishments	Households	Retail (Businesses)(Stores)
Busses	Institutional Bathrooms	Salons
Cafeterias <sup>2</sup>	Institutional Facilities	Schools <sup>2</sup>
Camp Grounds	Institutional Kitchens <sup>2</sup>	Ships
Cars	Isolation Wards	Spas
Churches	Kennels	Sports Stadiums
Classrooms	Kitchens	Supermarkets
Colleges	Laboratories	Tanning Spas
Commercial Kitchens <sup>2</sup>	Laundries	Tattoo Parlors
Convenience Stores	Locker Rooms	Taxis
Convention Centers	Lodging Establishments	Trailers
Crime Scenes	Manufacturing Facilities	Trains
Dairy Farms	Medical Offices <sup>3</sup>	Transportation Terminals
Dental Offices <sup>3</sup>	Medical Research Facilities <sup>3</sup>	(Turkey)(poultry)(beef)(swine) Farms
Dressing Rooms	Mobile Homes	Universities
Egg Processing Plants	Motels	Veterinary Clinics
Factories	Nursing Homes <sup>3</sup>	Wholesale Stores
Fast Food Operations	Office Buildings	Zoos
Federally Inspected Meat & Poultry Plants <sup>2</sup>		
Hatcheries(:)		
Egg Receiving Area	Tray Dumping Area	Chick Processing Area
Egg Holding Area	Chick Holding Area	Chick Loading Area
Setter Room	Hatchery Room	Poultry Buildings
Swine Premises(:)		
Farrowing (Barns)(Areas)	Hauling Equipment	Creep Area
Waterers	Dressing Plants	Chutes
Feeders	Blocks	Nursery

Bracketed information is denoted as: {insert as noted }, ( optional text ), [ required qualifier ] , <note>.  
The term "this product" used throughout this document may be replaced with the marketed product brand name.  
The term "x" is a place holder for numerical digits.  
Punctuation and plural/singular word forms may be adjusted to allow for grammatical correctness.  
Appropriate metric or imperial unit may be added as optional, supplemental information.



## **MICRO-ORGANISMS**

This product has been tested and found to be efficacious against the following micro-organisms on hard, non-porous surfaces:

<b>DISINFECTION</b>		
<b>Bacteria</b>		<b>Contact Time</b>
Pseudomonas aeruginosa	(ATCC 15442)	5 minutes
Staphylococcus aureus	(ATCC 6538)	5 minutes
Salmonella enterica	(ATCC 10708)	5 minutes
Enterococcus faecium	(ATCC 6569)	5 minutes
Escherichia coli	(ATCC 8739)	5 minutes
Escherichia coli O157:H7	(ATCC 43895)	5 minutes
Klebsiella pneumoniae	(ATCC 13883)	5 minutes
Listeria monocytogenes	(ATCC 984)	5 minutes
Proteus mirabilis (clinical isolate)		5 minutes
Proteus vulgaris	(ATCC 33420)	5 minutes
Salmonella (enteritidis) enterica	(ATCC 13076)	5 minutes
Salmonella (pullorum) enterica	(ATCC 9120)	5 minutes
Salmonella (typhi) enterica	(ATCC 6539)	5 minutes
Salmonella (typhimurium) enterica	(ATCC 14028)	5 minutes
Serratia marcescens (Industry Isolate)		5 minutes
Shigella flexneri	(ATCC 12022)	5 minutes
Shigella sonnei	(ATCC 9290)	5 minutes
Staphylococcus epidermidis	(ATCC 14990)	5 minutes
Streptococcus pyogenes	(ATCC 19615)	5 minutes
Yersinia enterocolitica	(ATCC 23715)	5 minutes
Antibiotic resistant Enterococcus faecium (clinical isolate) [Antibiotic Resistance to Ciprofloxacin, Penicillin, Vancomycin]		5 minutes
Antibiotic resistant Escherichia coli (clinical isolate) [Antibiotic Resistance to Methicillin]		5 minutes
Antibiotic resistant Klebsiella pneumonia (clinical isolate) [Antibiotic Resistance to Methicillin]		5 minutes
Antibiotic resistant Staphylococcus epidermidis (clinical isolate) [Antibiotic Resistance to Amoxicillin/CA, Ampicillin/Sulbatam, Cefazolin, Ciprofloxacin, Clindomycin, Erthromycin, Gentamicin, Levofloxacin, Oxacillin, Penicillin, Rifampin, Trimethsulfa]		5 minutes
Extended Spectrum Beta-Lactamase (ESBL) Escherichia coli [Antibiotic Resistance to Ceftazidime, Penicillin]	(ATCC BAA-196)	5 minutes
Methicillin resistant Staphylococcus aureus (MRSA)	(ATCC 33591)	5 minutes

Multi-Drug Resistant (MDR) Acinetobacter baumannii [Antibiotic Resistance to Amoxicillin/Clavulanic acid, Ampicillin/ Sublactam, Aztreonam, Benzylpenicillin, Cefepime, Cefotaxime, Cefoxitin, Ceftazadime, Ceftriaxone, Cefuroxime, Chloramphenicol, Clindamycin, Daptomycin, Erythromycin, Linezolid, Oxacillin, Penicillin, Quinupristin/Dalfopristin, Spectinomycin, Tetracycline, Tigecycline, Trimethoprim/sulfamethoxazole, Vancomycin ]	(ATCC 19606)	5 minutes
Oxacillin resistant Streptococcus pneumoniae	(ATCC 35088)	5 minutes
Vancomycin Resistant Enterococcus faecium (VRE) (clinical isolate) [Antibiotic Resistance to Ciprofloxacin, Penicillin, Vancomycin]		5 minutes
<b>Viruses</b>		<b>Contact Time</b>
Human Immunodeficiency Virus Type 1, (associated with AIDS)		30 seconds
Hepatitis B Virus (HBV) (Duck Hepatitis B Virus-DHBV)		3 minutes
Avian Influenza Virus (H9N2)		5 minutes
Bovine Viral Diarrhea Virus (BVDV)	(ATCC 534)	5 minutes
Hepatitis C Virus (HCV) (Bovine Viral Diarrhea Virus - BVDV)		5 minutes
Herpes Simplex Virus Type 1	(ATCC VR-733)	5 minutes
Herpes Simplex Virus Type 2	(ATCC VR-734)	5 minutes
Influenza Virus Type A2	(ATCC VR-544)	5 minutes
Pandemic 2009 H1N1 Influenza A Virus		
Pseudorabies	(ATCC VR-135)	5 minutes
Respiratory Syncytial Virus (RSV)	(ATCC VR-26)	5 minutes
Rotavirus	(ATCC VR-899)	5 minutes
SARS Associated Coronavirus		5 minutes
Vaccinia	(ATCC VR-119)	5 minutes
Vancomycin intermediate resistant Staphylococcus aureus (VISA)	(ATCC 700699)	5 minutes
<b>Fungi / Mildew</b>		<b>Contact Time</b>
Trichophyton mentagrophytes (athlete's foot fungus)	(ATCC 9533)	10 minutes
Aspergillus niger	(ATCC 6275)	10 minutes
Candida albicans	(ATCC 10231)	5 minutes
<b>SANITIZATION - NONFOOD</b>		
<b>Bacteria</b>		<b>Contact Time</b>
Staphylococcus aureus	(ATCC 6538)	15 Seconds
Klebsiella pneumoniae	(ATCC 4352)	15 Seconds
Enterobacter aerogenes	(ATCC 13048)	15 Seconds
(Pseudomonas aeruginosa	(ATCC 15442)	15 Seconds
Escherichia coli O157:H7	(ATCC 35150)	15 Seconds
Methicillin resistant Staphylococcus aureus (MRSA)	(ATCC 33592)	15 Seconds
Salmonella (typhi) enterica	(ATCC 6539)	15 Seconds
Streptococcus pyogenes	(ATCC 19615)	15 Seconds
Vancomycin resistant Enterococcus faecium (VRE)	(ATCC 51559)	15 Seconds

Bracketed information is denoted as: {insert as noted }, ( optional text ), [ required qualifier ] , <note>.  
The term "this product" used throughout this document may be replaced with the marketed product brand name.  
The term "x" is a place holder for numerical digits.  
Punctuation and plural/singular word forms may be adjusted to allow for grammatical correctness.  
Appropriate metric or imperial unit may be added as optional, supplemental information.

## **PRECAUTIONARY STATEMENTS**

### **PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS.**

#### **DANGER**

KEEP OUT OF REACH OF CHILDREN. CORROSIVE. Causes irreversible eye damage and skin burns. Do not get in eyes, on skin or on clothing. May be fatal if swallowed or inhaled. Do not breathe spray mist. Harmful if absorbed through skin. Wear goggles or face shield, rubber gloves and protective clothing when handling. Wash thoroughly with soap and water after handling and before eating, drinking, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

*<for containers 5 gallons or greater>*

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to fish, aquatic invertebrates, oysters and shrimp. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance contact your State Water Board or Regional Office of the EPA.

### **STORAGE AND DISPOSAL**

DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL.

**PESTICIDE STORAGE** - Store in a dry place no lower in temperature than 50°F or higher than 120°F.

**PESTICIDE DISPOSAL** - Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

<b>CONTAINER DISPOSAL</b> - <i>&lt;mandatory language - choose applicable directions&gt;</i>	
<i>For products only with household/residential use sites</i>	Nonrefillable container. Do not reuse this container. Triple rinse or equivalent. Then offer for recycling or reconditioning. If not available, dispose in trash
<i>For products with industrial, institutional, and commercial use sites</i>	Nonrefillable container. Do not reuse this container. Triple rinse or equivalent.
<i>For reuseable container</i>	Container can be reused once the container is empty. Empty container of concentrate into dilution dispenser (specific equipment can be named to replace "dilution dispenser"). Dispenser will automatically dilute a disinfectant solution. Remove concentrate label. Refill container with use-dilution of concentrate from the dispenser. Replace cap with spray nozzle provided. Place correct use dilution label on the spraybottle.
<i>For containers 5 gallons or less</i>	Triple rinse as follows: Fill container ¼ full with water and recap. Agitate vigorously. Follow Pesticide Disposal instruction for rinsate disposal. Drain for 10 seconds after the flow begins to drip. Repeat procedure two more times. Offer for

	recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill.
<i>For containers greater than 5 gallons</i>	Triple rinse as follows: Fill container ¼ full with water. Tip container on its side and roll it back and forth ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Offer for recycling or reconditioning, if available. If not available, puncture and dispose in a sanitary landfill.
<i>For products with industrial, institutional and commercial use sites and containers with a one-way opening for use in automatic dilution/dispersing systems</i>	Nonrefillable container. Do not reuse or refill this container. When container is empty, replace cap and discard in trash.  <b>Note to reviewer:</b> <i>The one-way opening container language is optional language and not required on the in-commerce label</i>

<when applicable>

**PACKET DISPOSAL** – Do not reuse; discard empty container and put in trash.

## **DIRECTIONS FOR USE**

*Note: product packaged in packet form must include Packet Directions for Use in addition to disinfection, sanitization, fungicide, cleaning, and/or site/application-specific directions for use.*

### **DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**DEODORIZATION:** To deodorize, apply this product as indicated under the heading DISINFECTION. For heavy duty deodorization dilute this product at 1 to 1.5 ounces per gallon and apply as directed under heading DISINFECTION.

**GENERAL CLEANING:** Apply this product to soiled area with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer. This product can be diluted at 0.5 ounces per gallon or other equivalent dilutions up to full strength. Thoroughly wet soiled surface and scrub as necessary.

**TO (CLEAN)(REMOVE) MOLD STAINS and MILDEW STAINS:** This product (product name) can be used at full strength or at a use dilution down to 0.5 ounces per gallon of water (or equivalent dilution). Apply full strength or diluted solution to surface to be cleaned with a cloth, mop, sponge or hand pump trigger sprayer. Thoroughly wet soiled surface completely and scrub as necessary.

**DISINFECTION:** To disinfect inanimate, hard non-porous surfaces add 0.5 ounces of this product per gallon of water. Apply solution with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow to remain wet for [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>] minutes, then remove excess liquid. For sprayer applications, spray 6-8 inches from surface, rub with brush, sponge or cloth. Do not breathe spray mist. For heavily soiled areas, a pre-cleaning step is required.

Prepare a fresh use-solution daily or more often if the use-solution becomes visibly (dirty), (contaminated) or diluted after use. All surfaces that come in contact with food must be rinsed with potable water. Do not use on utensils, glasses and dishes.

<Required if making an HIV claim on the product label>

**SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HCV and HBV OF SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS.**

**PERSONAL PROTECTION:** When handling items soiled with blood or body fluids, use disposable latex gloves, gowns, masks, and eye coverings.

**CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product.

**CONTACT TIME:** HIV-1 is inactivated after a thirty (30) second contact time. HBV is inactivated after a three (3) minute contact time. HCV is inactivated after five (5) minute contact time. Use either a five (5) or ten (10) minute contact time for other organisms as stated on the label.

**DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids must be autoclaved and disposed of according to local regulations for infectious waste disposal.

**SANITIZATION OF NON-FOOD CONTACT SURFACES:** To sanitize non-porous environmental surfaces add 0.5 ounces of this product per gallon of water. Apply solution with a mop, cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow to remain wet for 15 seconds, then remove excess liquid. For heavily soiled areas, a pre-cleaning step is required. This product cannot be used to clean utensils, glasses and dishes. Prepare a fresh solution for each use.

**FUNGICIDE & MILDEWCIDE:**

**Fungicide:** At 0.5 ounces per gallon dilution (equivalent dilution), this product is fungicidal against the pathogenic fungi, Trichophyton mentagrophytes (ATCC 9533) (Athlete's Foot Fungus) (a cause of ringworm) (a cause of ringworm of the foot) and Candida albicans (ATCC 10231). Apply this product (solution) with a cloth, sponge or hand pump trigger sprayer to hard, non-porous surface(s) (found in bathrooms, shower stalls, locker rooms, exercise facilities or other clean, non-food contact hard non-porous surfaces commonly contacted by bare feet). Allow the surface to remain wet for 10 minutes, for Candida albicans allow surface to remain wet for 5 minutes, then remove excess liquid. Diluted product must be applied daily or more frequently with heavy facility use.

**Mildewcide:** At 0.5 ounces per gallon (equivalent dilution) dilution, this product will kill Aspergillus niger (ATCC 6275) when applied to hard, non-porous surfaces. Apply this product (solution) with a cloth, sponge or hand pump trigger sprayer. Allow surface to remain wet for 10 minutes and let air dry. Repeat when mildew growth returns.

**MILDEWSTAT:** To control mold and mildew (Aspergillus niger) and the odors it causes on pre-cleaned, hard, non-porous non-food contact surfaces, add 0.5 (or 0.75) (or 1)(x) ounces of this product per gallon of water (or equivalent dilution). Apply solution with a cloth, mop, sponge or hand pump trigger sprayer making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth appears.

**SITE AND APPLICATION SPECIFIC DIRECTIONS FOR USE**

**Light Duty Cleaning (Directions):** Apply this product to soiled area with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer. This product can be diluted at 0.5 ounces per gallon or other dilutions up to 0.5 ounces per gallon. Thoroughly wet soiled surface and scrub as necessary.

**Disinfect Food Processing Premises (Directions):** For use on non-food contact, non-porous environmental surfaces such as floors, walls, bathrooms, office areas and storage areas, add 0.5 ounces of this product per gallon of water. For heavily soiled areas, a pre-cleaning step is required. Apply solution with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all

surfaces thoroughly. Allow to remain wet for [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>], then remove excess liquid. Before using this product, food products and packaging materials must be removed from the area or carefully protected.

**Restaurant Premises Disinfection (Directions):** For use on non-food contact, non-porous environmental surfaces such as floors, walls, bathrooms, office areas and storage areas, add 0.5 ounces of this product per gallon of water. For heavily soiled areas, a pre-cleaning step is required. Apply solution with a cloth, sponge or hand pump trigger sprayer so as to wet all surfaces thoroughly. Allow the surface to remain wet for [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>], then remove excess liquid. This product cannot be used to clean utensils, glassware and dishes.

**Food Contact Surfaces in Residential Kitchens:** To disinfect hard non-porous surfaces add 1 ounce of this product per gallon of water. Apply solution with a mop, cloth, sponge, hand pump trigger sprayer or low pressure coarse sprayer so as to wet all surfaces thoroughly. Allow to remain wet for [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>], then remove excess liquid. For sprayer applications, spray 6-8 inches from surface, rub with brush, sponge or cloth. Do not breathe spray mist. After use, food contact surfaces must be thoroughly rinsed with potable water. For heavily soiled areas, a pre-cleaning step is required.

Prepare a fresh use-solution daily or more often if the use-solution becomes visibly (dirty), (contaminated) or diluted after use. All surfaces that come in contact with food must be rinsed with potable water. This product cannot be used to clean utensils, glasses and dishes.

**Shoe Bath Sanitizer (Directions):** To prevent cross contamination into animal areas, and the packaging and storage areas of food plants, shoe baths containing one inch of freshly made use-solution should be placed at all entrances to buildings, hatcheries and all entrances to the production and packaging rooms. Scrape waterproof shoes and place in 0.5 ounces of this product per gallon of water (or equivalent use dilution) for 15 seconds prior to entering area. Change use-solution in the bath daily or sooner if use-solution appears dirty.

**Shoe Foam Sanitizer (Directions):** To prevent cross contamination into animal areas, and the packaging and storage areas of food plants, apply a foam layer approximately 0.5 to 2 inches thick made from a solution of 0.5 to 0.75 ounces per gallon of water (or equivalent use dilution) at all entrances to buildings, hatcheries, production and packaging rooms by using a foam generating machine or aerator to apply foam layer. Follow the foaming directions as specified by the manufacturer of the foam generator/aerator. Scrape waterproof shoes. Stand and/or walk through foamed area for 15 seconds prior to entering area. Foam area should be washed and replaced daily or when it appears dirty.

**To Control Mold and Mildew on Large, Inflatable, Non-Porous Plastic and Rubber Structures (animals, promotional items, moonwalk, slides, obstacle course play equipment, exercise equipment, and wrestling mats) (Directions):** Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 2.0 ounces of this product (product name) per 4 gallons of water (0.5 ounces per gallon of water) (or equivalent dilution) for a period of 10 minutes. Ventilate buildings and other closed spaces. Do not use equipment until treatment has been absorbed, set or dried.

**Veterinary Practice/Animal Care/Animal Laboratory/Zoos/Pet Shop/Kennels Disinfection (Directions):** For cleaning and disinfecting of the following hard non-porous surfaces: equipment not used for animal food or water, utensils, instruments, cages, kennels, stables, catteries, etc. Remove all animals and feed from premises, animal transportation vehicles, crates, etc. Remove all litter, droppings and manure from floors, walls and surfaces occupied or traversed by animals.

Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate surfaces with a use-solution of 0.5 ounces of this product per gallon of water (or equivalent dilution) for a period of [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>]. Ventilate buildings and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried.

**Farm Premise (Poultry) (Swine) (Beef) (Cattle) Disinfection (Directions):** Remove all animals and feed from premises, vehicles, coops, crates and enclosures. Do not use in milking stalls, milking parlors or milk houses. Remove all litter and manure from floors, walls and surfaces of barns, pens, stalls, chutes and other facilities and fixtures occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. For heavily soiled areas, a pre-cleaning step is required. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate all surfaces using a cloth, mop, or low pressure coarse sprayer to thoroughly wet surface or immerse all halters, ropes and other types of equipment used in handling and restraining animals as well as forks, shovels and scrapers used in removing litter and manure, with a use-solution of 0.5 ounces of this product per gallon of water (or equivalent dilution) for a period of [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>]. Ventilate buildings, cars, trucks, boats and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed, set or dried. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent and rinse with potable water before reuse. Prepare a fresh solution after each use.

**Hatcheries (Directions):** Use 0.5 ounces of this product per gallon of water to treat the following hard non-porous surfaces: hatching, setters, trays, racks, carts, sexing tables, delivery trucks and other hard surfaces. All treated surfaces exposed to the use-solution must remain wet for [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>] or more. Rinse with potable water before reuse and allow to air dry.

**Vehicles (Directions):** Clean all vehicles including mats, crates, cabs, and wheels with high pressure water. For heavily soiled vehicles use a detergent cleaner or this product at 0.5 or 0.75 ounces per gallon. Use 0.5 ounces per gallon of this product to treat all vehicles. All treated surfaces exposed to the use-solution must remain wet for [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>] or more. (Vehicle must air dry prior to entering or re-entering the roadway).

**Dressing Plant Use (Directions):** Disinfect equipment, utensils, walls and floors in poultry and animal dressing plants. Disinfect offal rooms, exterior walls and loading platforms of dressing plants. Cover or remove all food and packaging materials. Remove all gross soils. Saturate all surfaces with the use-solution: 0.5 ounces of this product per gallon of water (or equivalent dilution). Scrub to loosen all soils. Allow to soak for [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>] and thoroughly rinse all wetted and cleaned surfaces with potable water.

**Ultrasonic Bath Disinfectant (Directions):** Use this product to disinfect hard non-porous non-critical objects compatible with Ultrasonic cleaning units. Pour fresh solution at 0.5 ounces of this product per gallon of water directly into bath chamber. Pre-clean heavily soiled objects. Place objects into unit and operate for a minimum of [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>], according to the manufacturer's directions. Remove objects and wash with sterile water (sterile water for injection) or allow to air dry. Prepare fresh solution for each use.

**Disinfection of Hard Non-Porous Surfaces in Whirlpool Units (Directions):** After using the whirlpool unit, drain and refill with fresh water to just cover the intake valve. Add 0.5 ounces of this product for each gallon of fresh water added. Start the pump to circulate the solution. Wash down the unit sides, seat of the chair, lift and any/all related equipment with a clean swab, brush or

sponge. Treated surfaces must remain wet for [5 minutes][10 minutes *<if claiming Trichophyton mentagrophytes or aspergillus niger>*]. After the unit has been thoroughly cleaned, drain the solution from the unit and rinse any/all cleaned surfaces with fresh water. Repeat for heavy soiled units.

**Commercial Florist Use (Directions):** To clean, disinfect and deodorize hard non-porous surfaces in one step, prepare use-solution by adding (mixing) 0.5 ounces per gallon of water (or equivalent dilution) Remove all leaves, petals, garbage and refuse. Pre-clean surfaces using pressurized water where possible. Apply use-solution to hard (inanimate) non-porous surfaces thoroughly wetting surfaces as recommended and required, with a cloth, mop, brush, sponge or coarse sprayer. For heavily soiled areas, a preliminary cleaning is required. For sprayer applications, use a coarse mist pump or trigger sprayer. Spray 6-8 inches from surface; rub with brush, sponge or cloth. Do not breathe spray mist. Treated surfaces must remain wet for [5 minutes][10 minutes *<if claiming Trichophyton mentagrophytes or aspergillus niger>*]. Allow to air dry. Prepare a fresh solution at least daily or sooner if use-solution becomes visibly dirty.

**Disinfection of Barber/Salon Tools (Directions):** Immerse pre-cleaned barber/salon tools (combs, brushes, razors, manicure/pedicure tools, clippers, scissors, trimmer blades) in a use-solution (or equivalent dilution) of 0.5 ounces of this product per gallon of water for 5 (to 10) minutes. Rinse instruments thoroughly and dry before reuse. A fresh use-solution must be prepared daily or more often if the use-solution becomes (visibly dirty) (contaminated) (cloudy). **NOTE:** Plastics may remain immersed until ready to use. Stainless steel shears and instruments must be removed from the use-solution after [5 minutes][10 minutes *<if claiming Trichophyton mentagrophytes or aspergillus niger>*], rinsed, dried and kept in a clean, non-contaminated receptacle. Prolonged immersion may cause damage to stainless steel or metal instruments.

*<Not for use in California>*

**ENTRYWAY SANITIZATION SYSTEMS (Directions):** To prevent cross contamination from area to area, set the system to deliver (0.5 to 0.75 ounces per gallon of water) (or equivalent use dilution) of sanitizing solution. The (spray) (foam) should cover the entire path of the doorway. Set the system so that a continuous wet blanket of sanitizer solution is delivered to the floor. Do not mix other additives to the sanitizing solution.

**WATER AND SMOKE DAMAGE RESTORATION AGAINST ODOR-CAUSING BACTERIA AND FUNGI FOR INSTITUTIONAL, INDUSTRIAL AND HOSPITAL USE (Directions):**

this product is suitable for use in water damage restoration situations against odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushion, sub floors, drywall, trim, and frame lumber, tackless strip and paneling. Mix 0.5 ounces of this product per gallon of water, saturate affected materials with enough use-solution to remain wet for at least 5 (to 10) minutes. Use proper ventilation.

**SEWER BACKUP & RIVER FLOODING (Directions):** During mitigation procedures, dilute 0.5 to 1.0 ounces of this product per gallon of water allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil along with non-salvageable materials. Saturate all affected areas with a sprayer using a coarse spray tip, allow material to remain wet for 10 minutes before cleaning and extraction.

*(Not for use in California)*

**CARPETS, CARPET CUSHIONS AND OTHER POROUS MATERIALS SUCH AS SUB FLOORS, DRYWALL, TRIM, AND FRAME LUMBER, TACKLESS STRIP AND PANELING (Directions):**

For water damage from a clean water source, extract excess water. Test hidden area for colorfastness. Mix 0.5 to 1.0 ounces of this product per gallon of water, allowing for the diluting effect of absorbed water within saturated materials. Remove gross filth or heavy soil. Apply directly with a sprayer using a coarse spray tip, to fully saturate affected materials. Roll, brush or agitate into materials and allow the materials to remain damp for [5 minutes][10 minutes *<if claiming*



*Trichophyton mentagrophytes* or *aspergillus niger*>]. Follow with a thorough extraction. Dry rapidly and thoroughly.

**CARPET DEODORIZER AGAINST ODOR-CAUSING BACTERIA FOR INSTITUTIONAL, INDUSTRIAL AND HOSPITAL USE (Directions):** This product deodorizes the carpet by controlling/reducing the growth of odor-causing bacteria. It can be used in industrial and institutional areas: motels, hotels, nursing homes and hospitals. Vacuum carpet thoroughly prior to application.

Mix 0.5 ounces of this product per gallon of water. Follow the Injection and/or Extraction procedures as specified for any conventional potable steam cleaning equipment you are using. For rotary floor machines, mix 1 ounce per gallon of water and spray on carpet at a rate of 300-500 sq. ft. per gallon. For truck mounted extraction machines mix 12 ounces of this product per gallon of water and meter at 4 gallon per hour.

For use on washable synthetic fibers. Do not use on wool. Test colorfastness of carpet before use. Apply diluted product to a small concealed spot, then rub with a clean white cloth. If color changes or transfers to cloth, a water-based product should not be used.

After using the product, set carpet pile in one direction with a stiff brush. Place aluminum foil under the legs of furniture while carpet is drying. Over-wetting can cause carpet to shrink. Manufacturer assumes no responsibility for over-wetting misuse.

**NOTE:** This product must not be mixed with other cleaning products. Follow the cleaning procedures as specified by the manufacturer of the cleaning equipment.

**MUSHROOM FARM SANITIZATION** – This product disinfects, cleans, and deodorizes in one step. Apply to surfaces which do not come in contact with the crop, compost or casing. Remove gross filth and heavy soil deposits using a shovel, broom or vacuum depending on the area to be disinfected. For heavily soiled areas, a pre-cleaning step is required. Apply this product at 0.5 ounces per gallon of water dilution, thoroughly wetting the surfaces with a mop, sponge, cloth, spraying or soaking. Treated surfaces must remain wet for [5 minutes][10 minutes <if claiming *Trichophyton mentagrophytes* or *aspergillus niger*>] and let air dry. The use-solution is prepared fresh for each use, then discarded. If greater detergency is required, dilute this product at 1 to 2 ounces per gallon of water. Bactericidal when diluted at 0.5 ounces per gallon, this product demonstrates effective disinfectant activity against the following **mushroom** pathogens: *Pseudomonas tolaasii* (Blotch Disease) and *Pseudomonas* sp. (Mummy Disease). **DO NOT APPLY TO MUSHROOM CROP, COMPOST OR CASING.** For Foot Dip of Waterproof Footwear use this product at 0.5 ounces per gallon of water in foot dip tray. Shoe baths must contain at least 1 inch of freshly made solution and must be placed at the entrances to buildings. Scrape shoes and place in diluted solution for 5 (to 10) minutes before entering building. Change solution in bath daily or sooner if solution appears dirty.

**CITRUS CANKER CONTROL** – For prevention of Citrus Canker Disease through treatment of equipment. This product, when used at a 1:164 dilution (2000 ppm active quaternary) (or) (0.80 ounces per gallon of water) (or equivalent dilution), is effective against *Xanthomonas axonopodis* pathovar citri (Citrus Canker). Treat all trucks, vehicles, and equipment thoroughly at the 1:164 dilution. Treatment can be applied to trucks, attached trailers, field harvesting equipment; including cargo area, wheels, tires, under carriage, hood, roof, fenders and any other part of transportation equipment that can be taken into infested areas. To prevent the spread of Citrus Canker by this artificial means of transportation, treatments are made by trigger spraying, dipping or brushing, using the maximum rate of 2000 ppm quaternary. After use, all surfaces which come in contact with food or crop must be rinsed with potable water.

### **DIRECTIONS FOR FOGGING DEVICES**

**Directions for Hatchery Rooms Using Fogging Devices:** Remove all animals and feed from premises, vehicles and enclosures. Remove all litter and manure from floors, walls and surfaces of the room to be treated. Empty all troughs, racks and other feeding and watering appliances. Close room off so fog is confined to room to be treated. Wear a dust mist respirator when mixing the use-solution and pouring it into the fogging apparatus. Mix .25 parts of this product to one and a quarter parts water (0.64 gallons of this product to 5 gallons water). Insert the nozzle of the fogger through a suitable opening into the room. With the setting in maximum output, fog for one minute for each 4000 cubic feet of space in the room. When fogging is completed, ventilate buildings and other closed spaces. Do not house livestock or employ equipment until treatment has been absorbed or dried. Thoroughly clean all surfaces with soap or detergent and rinse with water.

**Directions for Incubators and Hatchers using Fogging Devices:** Wear a dust mist respirator when mixing the use-solution and pouring it into the fogging apparatus. Mix 3.85 ounces of this product to 122 ounces of water. Fog 0.75- 2 ounces of this use-solution into setters and hatchers immediately after transfer. Repeat daily in setters and every 12 hours in hatchers. Discontinue hatcher treatments at least 24 hours prior to pulling the hatch. Do not allow people to contact or breathe this fog and do not enter until the fog has settled (30-60 minutes after fogging is completed). It is acceptable to fog setters and hatchers with a 0.5 ounce per gallon use-solution of this product on an hourly or every other hour basis. If this is done, fog for 30-90 seconds once per hour or once every two hours. Thoroughly scrub all treated feed racks, mangers, troughs, automatic feeders, fountains and waterers with soap or detergent, and rinse with potable water before reuse.

**Directions for Water and Smoke Damage Restoration Using Fogging Devices:** This product is suitable for use in water damage restoration situations to control odor causing bacteria and fungi. Clean the room to be treated to remove all litter and gross filth from floors, walls and surfaces. Close room off so fog is confined to room to be treated. Wear a dust mist respirator when mixing the use-solution and pouring it into the fogging apparatus. Mix 0.5 parts of this product to 0.5 parts water (0.5 gallons of this product to 0.5 gallons water). Insert the nozzle of the fogger through a suitable opening into the room. With the setting in maximum output, fog for one minute for each 4000 cubic feet of space in the room. When fogging is completed, ventilate buildings and other closed spaces.

**NOTE:** The fog generated is irritating to the eyes, skin and mucous membranes. Do not under any circumstances enter a room or building within two hours of the actual fogging and a minimum of 4 air exchanges (ACH) per hour in the facility. If the building must be entered, the individuals must wear a self-contained respirator approved byNIOSH/MSHA, goggles, long sleeves and long pants.

## **DILUTION SYSTEM SPECIFIC DIRECTIONS FOR USE**

**(Automatic Dilution)(Dilution System Name) ((Closed)(Open) Loop):** (Concentrated product – dilute before using)

ATTACH this product (PRODUCT NAME) TO THE AUTOMATIC DILUTION CENTER: Remove white cap from product bottle. DO NOT REMOVE purple PLUG FROM BOTTLE THROAT! Attach the bottle to the automatic dilution center by screwing on matching purple dispensing cap connected to dilution center. Push down while screwing on cap. This dispensing system delivers 0.5 oz. per gallon of water (1:256) use solution.

**Automated Dilution System (Dilution System Name):** Remove cap/spray nozzle from empty container. Fill empty container with a newly prepared use-solution. Replace cap/spray nozzle. Place correct use-dilution label on newly filled container.

### **Smart Mix Pro Dispensing System:**

1. Fill Smart Mix Pro pouch with cold water to the fill line arrow below the cap.
2. Fold and Squeeze the chemical concentrate to burst inner pouch.
3. Shake the bag to mix chemical fully.
4. Use the push button to fill the bottle.

Once water and concentrate are mixed, 1 gallon (3.8 Liters) of ready to use (disinfecting/cleaning) (disinfecting) solution.

### **Integrated Dispensing System (Name of Integrated Dispensing System) (Control Knob Key) -**

Attach to source of potable water with quick connect attachment.



Rotate control knob to (1:256)( 1:256 ) for low flow rate applications such as filling trigger spray bottle.



Rotate control knob to (1:256)( 1:256 ) for high flow rate applications as filling a mop bucket.

Squeeze handle to dispense a 0.5 oz. per gallon of water (1:256) use solution of product.

**PACKETS:** *Pre-measured tear open packets* – (Simply) (tear) open and pour contents into X gallons of water (for a 1:256 dilution). Keep packets in box until ready to use.

or

Pour contents of (x oz)(0.5 or 1 oz) packet into (x)(1 or 2) gallons of water to yield 1:256 dilution. Keep packets in box until ready to use.

**The following language is required for products supplied with supplemental labeling: inserts, hang tags and reference sheets.**

**The following language is required to appear on inserts, hangtags and reference sheets:**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Labeling must be in the possession of the user at the time of using the product. See container label for FIRST AID and PRECAUTIONARY STATEMENTS and complete DIRECTIONS for USE before applying. Use of this product is subject to the precautions and limitations imposed by the label affixed to the container for this product.

**The following language is required to appear on the container label:**

See (insert) (hangtag) (reference sheet) for (complete) (additional) information.

## **SECONDARY / SERVICE CONTAINER LABELING**

**<Secondary / service container labeling for use only when this product is diluted in accordance with 0.5 ounces per gallon (or equivalent dilution). Not for use with alternate dilutions.>**

*When this product is diluted in accordance with the directions on this label, the dilution container must bear the following statements:*

### **SC-5:256N / Product Brand Name 1839-235**

Didecyl dimethyl ammonium chloride .....	19.596%
Alkyl (50% C <sub>14</sub> , 40% C <sub>12</sub> , 10% C <sub>16</sub> )....	13.064%
dimethyl benzyl ammonium chloride	
<b>OTHER INGREDIENTS .....</b>	<b>67.34%</b>
<b>Total.....</b>	<b>100.000%</b>

*The product in this container is diluted as directed on the pesticide product label. Follow the directions for use on the pesticide label when applying this product. Further sale or distribution of this product is strictly prohibited.*

### **PRECAUTIONARY STATEMENTS**

**KEEP OUT OF REACH OF CHILDREN.** Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Harmful if absorbed through skin. Wash thoroughly with soap and water after handling and before eating, drinking, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse.

**PACKET LABEL UNIT PACKAGING**

**(0.5, 1, or X) OUNCE PACKET LABEL TO BE USED WITH MASTER CONTAINER LABEL:  
MIX EACH PACKET WITH (2, 4, or X) GALLONS OF WATER:**

*(Note to reviewer: X ounce packets and corresponding X amount of water will be adjusted to yield equivalent dilution in accordance with directions for use on this label)*

**SC-5:256N / Product Brand Name  
1839-235**

Didecyl dimethyl ammonium chloride ..... 19.596%  
Alkyl (50% C<sub>14</sub>, 40% C<sub>12</sub>, 10% C<sub>16</sub>).... 13.064%  
dimethyl benzyl ammonium chloride  
**OTHER INGREDIENTS** .....67.34%  
**Total**.....100.000%

**KEEP OUT OF REACH OF CHILDREN  
DANGER**

See master label for additional precautionary statements and use directions.  
**Disposal** – Do not reuse; discard empty packet by putting in trash.

EPA REG. NO. 1839-235 EPA EST. NO. (1839-IL-01)(xxxx-xx-xxx)  
STEPAN COMPANY  
22 West Frontage Road • Northfield, IL 60093  
Pour contents of packet into XX gallons of water for a 1:256 dilution.  
NET CONTENTS: XX (ounces)(oz) per packet

## GRAPHICS

### Automatic Dilution System



### Integrated Dispensing System



### Smart Mix Pro™ dispensing system

