

This is a concentrated insecticide and must be diluted for use. Each gallon of concentrate contains 4 pounds of Malathion (1/2 pound per pint). Mix by adding to water in spray tank with pump and agitator running.

FLIES To control flies around animal pens, coops, barns and around homes, mix 1 teaspoon concentrate per pint of water. Apply to thoroughly wet surfaces, stanchions, floors, windows in dairy barns, fences, around garbage cans, etc. For bait spray add 1/2 tablespoon sugar or 2 teaspoons of corn syrup or unsulfurized molasses per gallon of above solution. Apply as above or as spray by drenching over surface of manure. In loafing sheds, spray the dry bedding within 18 inches of walls and upright bracings. Do not use in milk rooms. Remove animals from building before treating.

Do not use in dwellings.
Do not use in poultry houses.

In conjunction with the malathion treatments in and around dairy barns and other agricultural premises, fly breeding sites such as manure and other waste material, should be eliminated.

DOGS AND CATS—For the control of fleas on dogs and cats, apply a spray containing 1/2 teaspoon of malathion 50% emulsifiable liquid per 1 pint of water. Wet thoroughly by means of a hand sprayer.

If the animal has free run, treatment may have to be made every second or third week. To avoid unnecessary applications, animal should be closely examined for fleas before re-treating. If the animal is confined to an area free of fleas, one treatment may be sufficient.

TRIANGLE

QUALITY TESTED

**50% MALATHION
PREMIUM GRADE**

Emulsifiable Concentrate

ACTIVE INGREDIENTS:

Malathion*	50%
Petroleum Hydrocarbons	45%
INERT INGREDIENTS	05%
	100%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

**CAUTIONS: Keep Out of Reach of Children.
See additional cautions in side panel.**

Net Contents
1 PINT

FLEAS IN ANIMAL QUARTERS—For the control of fleas in dog and cat kennels, in pens, yards, lawns, and under houses, use a spray containing 2/3 cup of 50% emulsifiable liquid per 1 gallon of water at the rate of 1 gallon per 1,000 sq. ft.

Use a garden sprayer. Remove manure or debris before spraying.

ORNAMENTALS—For control of Aphids, Spider Mites, White Fly, Mealy Bugs, Thrips, Japanese Beetle, Adult Four Lined Bug, Tarnished Plant Bug, Rose Leafhopper and Potato Leafhopper on ornamentals apply 2 pints in 100 gallons of water (2 teaspoonfuls per gal.). Make thorough full coverage sprays with repeated applications as needed. Injury may occur on certain ferns, including Boston, Maidenhair and Peteris as well as some species of Crassola.

CAUTIONS

Use only as directed. Harmful if inhaled, swallowed or absorbed through the skin. Avoid breathing mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after using. Avoid contamination of feed and food stuffs. Keep out of reach of children and domestic animals. Malathion, upon repeated or prolonged careless use, may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid. Atropine is antidotal. "This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label."

USDA Reg. No. 1842-69

MANUFACTURED BY
TRIANGLE CHEMICAL CO.
MACON GEORGIA

ACCEPTED

Dec. 9, 1969
IN THE UNITED STATES OF AMERICA
FOR THE FEDERAL BUREAU OF INVESTIGATION
ED. W. ...
TO ALLIANCE ...

water in spray tank with pump and
ing.

control flies around animal pens,
and around homes, mix 1 teaspoon
per pint of water. Apply to thoroughly
stanchions, floors, windows in dairy
around garbage cans, etc. For bait
tablespoon sugar or 2 teaspoons of
unsulfurized molasses per gallon of
n. Apply as above or as spray by
er surface of manure. In loafing sheds,
bedding within 18 inches of walls and
ngs. Do not use in milk rooms. Re-
from building before treating.
in dwellings.
in poultry houses.

tion with the malathion treatments in
dairy barns and other agricultural
breeding sites such as manure and
material, should be eliminated.

D CATS--For the control of fleas on
ts, apply a spray containing 1/2 tea-
malathion 50% emulsifiable liquid per
ater. Wet thoroughly by means of a

al has free run, treatment may have to
y second or third week. To avoid un-
lications, animal should be closely ex-
leas before re-treating. If the animal
an area free of fleas, one treatment
ient.

TRIANGLE

QUALITY TESTED

50% MALATHION PREMIUM GRADE

Emulsifiable Concentrate

ACTIVE INGREDIENTS:

Malathion*	50%
Petroleum Hydrocarbons	45%
INERT INGREDIENTS	05%
	<hr/> 100%

*O'-O-dimethyl dithiophosphate of diethyl mercaptosuccinate

**CAUTIONS: Keep Out of Reach of Children.
See additional cautions in side panel.**

lawns, and under houses, use a
2/3 cup of 50% emulsifiable liqu
of water at the rate of 1 gallon

Use a garden sprayer. Remove i
before spraying.

ORNAMENTALS--For control o
Mites, White Fly, Mealy Bugs,
Beetle, Adult Four Lined Bug,
Bug, Rose Leafhopper and Potato
ornamentals apply 2 pints in 100
(2 teaspoonfuls per gal.). Make tho
age sprays with repeated applicati
jury may occur on certain ferns, i
Maidenhair and Peteris as well as
Crassola.

CAUTIONS

Use only as directed. Harmful if i
ed or absorbed through the skin.
mist. Avoid contact with skin, e
Wash thoroughly with soap and w
Avoid contamination of feed and f
out of reach of children and do
Malathion, upon repeated or pronon
may cause cholinesterase inhibition.
curs, get prompt medical aid. Atrop
"This product is toxic to fish. K
body of water. Apply this produc
fied on this label."

USDA R

MANUFACTURED BY
TRIANGLE CHEMICAL CO.
MACON, GEORGIA

ACCEPTED

Dec. 9, 1969

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ANTICIDE ACT
FOR ECONOMIC PEST CONTROL REGISTERED
ED UNDER NO. 184269 SUBJECT
TO ATTACHED COMMENTS.