## 15% FERMATE DUST

A Ferbam fungicide for control of Blue Mold on tobacco plant beds.

### **ACTIVE INGREDIENTS:**

\*Ferbam (Ferric dimethyl dithiocarbamate

11.4%

INERT INGREDIENTS

88.6°

100.00%

\*Equivalent to 15% Fermate.

CAUTION: Keep Out of Reach of Children.

#### DIRECTIONS FOR USE

Blue Mold in tobecco plant beda. Apply directly to the young plants or dust through the plant bed cloth. Apply 2 to 3 pounds of dust per plant bed (100 Sq. Yds.) depending on size of plants. Start application when leaves are about size of a dime, or when Blue Mold first appears in the vicinity. Apply at 3 to 4 day intervals until transplanting time.

#### **CAUTIONS**

MAY CAUSE IRRITATION OF EYES, NOSE, THROAT, AND SKIN.

Avoid breathing dust or spray mist.

Avoid contact with skin, eyes, and clothing.

Keep away from fire or sparks.

In case of contact, flush with plenty of water: for eyes, get medical attention.

Do not reuse container. Destroy by burying or burning.

Stay out of smoke and functionally

Seller warrants that this product conforms to it chemical description and is reasonably fit for th purpose stated on the label when used in accordance with directions under normal conditions of use, bu neither this warranty nor any other warranty of marchentability or fitness for a particular purpose cupressed or implied, extends to the use of this product contrary to abel instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the seller, and buyer assumes the risk of any such

Manufactured by

# TRIANGLE CHEMICAL COMPANY MACON, GA.

Net Wt.

Lbs.

USDA Reg. No. 1842-125

ACCEPTED

MAY 2 9 1968

UNDER THE TANAL INSECTIONS
OF UNGICIDE AND HODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED UNDER NO. 13 72-16 SUBJECT
TO ATTACHED COMMENTS.

1847-105