

PM 16 1792-130 1/2
12-31-89

**PRECAUTIONARY STATEMENTS
WARNING!**

Use only as directed. Harmful if inhaled, swallowed, or absorbed through the skin. Avoid breathing mist. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after using. Avoid contamination of feed and food stuffs. Keep out of reach of children and domestic animals. Malathion, upon repeated use, may cause cholinesterase inhibition. If poisoning occurs, get prompt medical aid.

NOTE TO PHYSICIAN

Malathion may cause cholinesterase inhibition. ATROPINE is antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

DISPOSAL

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

REV A 86060 S



**57% MALATHION
PREMIUM GRADE
EMULSIFIABLE CONCENTRATE**

ACTIVE INGREDIENTS

Malathion*	57%
Xylene	38%
INERT INGREDIENTS	05%
	100%

(*2,0-dimethyl dithiophosphate of diethyl mercaptosuccinate)

WARNING!

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Mfg. By
TRIANGLE CHEMICAL CO
Macon, Georgia

EPA Reg. No. 1842-132
EPA Est. No. 1842-GA-1

NET CONTENTS... 1 GALLON

Do not treat animals under 1 month of age.

If the animal has free run, treatments may have to be made every second or third week. To avoid unnecessary applications, animal should be closely examined for fleas before re-treating. If the animal is confined to an area free of fleas, one treatment may be sufficient.

FLEAS IN ANIMAL QUARTERS - For the control of fleas in dog and cat kennels, in pens, yards, lawns, and under houses, use a spray containing 1/2 cup of 57% emulsifiable liquid per 1 gallon of water at the rate of 1 gallon per 1,000 square feet.

Use a garden sprayer. Remove manure or debris before spraying.

ORNAMENTALS - For control of Aphids, Spider Mites, White Fly, Mealy Bugs, Thrips, Japanese Beetle Adult, Four Line Leaf Bug, Tarnished Plant Bug, Rose Leafhopper and Potato Leafhopper on ornamental apply 2 pints in 100 gallons of water; 2 cap. per gallon. Make thorough full coverage sprays with repeated applications as needed. Injury may occur on certain ferns, including Boston, Maidenhair and Peteris as well as some species of Crassula.

NOTICE OF WARRANTY

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the seller, and buyer assumes the risk of any such use.

DIRECTIONS FOR USE

This is a concentrated insecticide and must be diluted for use. Each gallon of concentrate contains 5 pounds of Malathion. Mix by adding to water in spray tank with pump and agitator running.

FLEES - To control flies around animal pens, coops, barns and around homes, mix 1/2 cup concentrate per gallon of water. (2 gallons to 100 gallons.) Apply to thoroughly wet surfaces where flies light or congregate, such as walls, ceilings, stanchions, floors, windows in dairy barns, fences, around garbage cans, etc. For bait spray add 1/2 cup sugar or 1/2 cup corn syrup or unsulfurized molasses per gallon of above solution. Apply as above or as a spot treatment. For fly maggots in manure, apply bait spray by drenching over surface of manure. In loafing sheds, spray the dry bedding within 18 inches of walls and upright bracing. Do not use in milk rooms. Remove animals from building before treating.

NOT FOR USE IN POULTRY HOUSES. DO NOT USE IN DWELLINGS. In conjunction with the malathion treatment in and around dairy barns and other agricultural premises, fly breeding sites such as manure and other waste material, should be eliminated.

DOGS & CATS - For the control of fleas on dogs and cats, apply a spray containing 2 tbs. of malathion 57% emulsifiable liquid per 1 gallon of water. Wet thoroughly by means of a hand sprayer. Do not spray directly on dogs or cats.

BEST AVAILABLE COPY

Not Reviewed. Registrant claims to be in accordance with
Reregistration of Pesticide Product Guidance for Phase 2
Response, Page 2.12, Active to Inert Change in Status.