

Endrin-Methyl Parathion Emulsifiable Concentrate

6/17/70
 1824-757

For use primarily with ground rigs. If airplane use is desired, use at least 3 to 10 gals water per acre.

This is a concentrated insecticide and must be diluted with water for use. Mix by adding to spray tank containing water with pump and agitator running. Before using study safety directions. Consult your physician about obtaining an adequate supply of 1/100 gr. Atropine tablets for emergency use.

DIRECTIONS FOR USE

FOR COFFEE: bollweevil, bollworm, thrips, fleahoppers, cutworms, grasshoppers, aphids, apply 1 to 2 pints per acre (depending on size of plants) at 3 to 7 day intervals or as necessary.

In the application of this concentrate, use the following directions:

1. Do not spray on windward side of field.

2. Do not spray on treated plants within 24 hours of application. Should wear protective clothing, do not handle or handle clothing of applicator. Do not use on or near animals. Do not spray on or near treated fields. Do not spray or allow drift to land on crops, ornamentals, trees, shrubs, or other plants.

3. Do not use to fish and wildlife. Do not contaminate body of water nor apply to any body of water. See the label. Birds feeding on treated plants may be killed. Do not apply when wind is strong or drift from treated area. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of water.

Contains 1.6# Endrin plus 1.6# Methyl Parathion per gallon

ACTIVE INGREDIENTS:

*Endrin	18.3%
**O,O-dimethyl O-p-nitrophenyl phosphorothioate	18.3%
Xylene	57.4%
INERT INGREDIENTS	6.0%
	100.0%

*hexachlorocyclohexane, endo dimethanonaphthalene **Methyl Parathion

DANGER: Keep Out of Reach of Children.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME

DANGER

Poisonous by skin contact inhalation, or swallowing. Rapidly absorbed through skin. Repeated exposure may without symptoms, be increasingly hazardous.

DO NOT GET ON SKIN ON CLOTHING OR IN EYES

Wear heavy natural rubber gloves and clean, waterproof or freshly laundered protective clothing (coveralls, cap, etc.) and goggles. Destroy and replace gloves frequently. Wash thoroughly with soap and warm water before smoking or eating. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and water before reuse. In case of contact, immediately remove contaminated clothing and wash thoroughly with soap and warm water. In case of eye contact, flush with plenty of water.

DO NOT BREATHE DUST, VAPOR OR SPRAY MIST

Use respirator of type passed by the American Manufacturer Methyl Parathion and Endrin protection. Airplane pilots should wear full face mask of type marked for harmful vapors, provide aircraft with full ventilation. Keep all persons out of operating areas or vicinity where there may be danger of poisonous drift. Vacate area immediately after treatment and do not re-enter until area has been sprayed.

DO NOT CONTAMINATE FOOD OR FOOD PRODUCTS

Bury spillage. Clean up area with strong lye solution. Burn or bury containers. Never re-use. Stay out of smoke or fumes.

PHYSICIANS NOTE

Treat Symptomatically

Warning symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin point pupils, salivating, nausea, vomiting, diarrhea and cramps.

Methyl Parathion is a cholinesterase inhibitor and treatment of poisoning should include the use of atropine in large dose to produce full atropinization and artificial respiration, if necessary. Endrin paralyses central nervous system, convulsions which responds rather well to almost any of the barbituric acid derivatives. The clinical acute period following Methyl Parathion poisoning lasts 24 to 48 hours. Acute toxic effects of both poisons might well last several days if a weak or naive.

Repeated exposure to cholinesterase inhibitors may without warning cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until full cholinesterase regeneration has been established as determined by blood test.



Atropine is an antidote. Consult physician for emergency supply.

ANTIDOTE

Call a physician at once in all cases of suspected Methyl Parathion poisoning. If symptoms or signs of poisoning include blurring vision, abdominal cramps and tightness in the eyes, do not wait for doctor's diagnosis. Give atropine tablets until the gums are dry. Remove victim from area of exposure. If Methyl Parathion is present, remove contaminated clothing and wash skin thoroughly with soap and water. In case of ingestion of Methyl Parathion, induce vomiting if possible. Do not give anything to eat or drink. If patient is unconscious, do not give anything by mouth. If patient is conscious, give small sips of water. If patient is unconscious, give small sips of water. If patient is unconscious, give small sips of water.

NOTICE OF WARRANTY

This product is warranted to be effective under normal conditions of use. It is not warranted to be effective under conditions of use not specified on the label. The warranty does not extend to any other use of the product. The warranty does not extend to any other use of the product. The warranty does not extend to any other use of the product.

Manufactured by

TRIANGLE CHEMICAL COMPANY

MACON, GA