

# 10-1 SEVIN PARATHION DUST

FOR USE ON LEEA MO TOMATO AND PEPPER INSECTS

## ACTIVE INGREDIENTS

Carbaryl 14 naphthyl 2,4-dimethyl 4-phenyl

Parathion 0.0001% (by weight) of the product

## INERT INGREDIENTS

ANGEL

DANGER



POISON

1-1-10-1

1842-206

DIRECTIONS FOR USE

# DANGER

y skin contact, inhala-  
wing. Rapidly absorbed  
in. Repeated exposure  
ymptoms, be increasing-  
Keep all persons out of  
vicinity where there  
of poisonous drift. Va-  
ould not be re-entered  
on residues have dis-

GET ON SKIN OR  
OR IN EYES.

1 rubber gloves and gog-  
waterproof or freshly  
ective clothing (cover-  
. Destroy and replace  
tly. Wash thoroughly  
d warm water before  
ing. Bathe immediately  
d change all clothing.  
; thoroughly with soap  
re re-use. In case of con-  
ely remove contaminated  
wash thoroughly with  
m water.

BREATH DUST, VAP-  
Y MIST.

Wear a mask or respirator of type  
passed by U. S. Department of Agri-  
culture for Parathion protection. Air-  
plane pilots should wear full face can-  
ister type mask.

If handled indoors provide mechan-  
ical exhaust ventilation.

**DO NOT CONTAMINATE FOOD  
OR FEED PRODUCTS.** Bury spil-  
lage. Clean up area with strong lye  
solution. Burn or bury containers. Nev-  
er re-use. Stay out of smoke or fumes.

## PHYSICIANS NOTE

*Warning symptoms include weakness, head-  
ache, tightness in chest, blurring vision, non-  
reactive pin point pupils, salivating, sweating,  
nausea, vomiting, diarrhea and abdominal  
cramps.*

*TREATMENT: Give atropine, preferably  
by injection, Grain 1/100, two or three tab-  
lets at once, and parenterally or orally every  
hour as required up to 30 tablets or until  
pupils dilate. Never give morphine. Clear  
chest by postural drainage. Artificial respira-  
tion or oxygen administration may be neces-  
sary. Observe patient continuously 48 hours.  
Repeated exposure to cholinesterase inhibi-  
tors may without warning cause prolonged  
susceptibility to very small doses of any  
cholinesterase inhibitor. Allow no further ex-  
posure until time for cholinesterase regenera-  
tion has been allowed as determined by blood  
tests.*

Atropine is an antidote. Consult  
physician for emergency supply.

## ANTIDOTE

Call a physician at once in all cases  
of suspected Parathion poisoning. If  
symptoms or signs of poisoning include  
blurred vision, abdominal cramps and  
tightness of chest don't wait for a  
doctor but give two atropine tablets  
(each 1/100 gr.) at once. Remove  
patient immediately from area where  
**Parathion is present. Remove contami-  
nated clothing and wash skin clean  
with plenty of soap and water, to re-  
move all traces of Parathion. If swal-  
lowed, induce vomiting by giving warm  
salty or soapy water and repeat until  
vomit fluid is clear. Do not give  
liquids to unconscious patient. Have  
patient lie down and keep quiet.**

## NOTICE OF WARRANTY

Seller warrants that this product conforms to its  
chemical description and is reasonably fit for the  
purpose stated on the label when used in accordance  
with directions under normal conditions of use, but  
neither this warranty nor any other warranty of  
merchantability or fitness for a particular purpose,  
expressed or implied, extends to the use of this  
product contrary to label instructions, or under  
abnormal conditions, or under conditions not rea-  
sonably foreseeable to the seller, and buyer assumes  
the risk of any such use.

USDA Reg. No. 1842-206

**IGLE CHEMICAL COMPANY**

**Macon, Georgia**