

5-1 SEVIN® PARATHION DUST

DIRECTIONS FOR USE

Note Cautions and read directions carefully before using.

For TOBACCO, APHIS, FLEA BEETLES, HORNWORMS.

Apply 15 pounds per acre as soon as plants become established in field. Repeat at two week intervals, increasing treatment to 20 pounds per acre by the time crop is half grown.

Do not apply within 5 days of pruning or 15 days of cutting. Avoid having plant pieces coming into contact with the skin.

TOMATOES AND PEPPERS, HORNWORMS, FRUITWORMS, APHIS, and LEAFMINER. When insects appear apply 20 to 30 pounds per acre, depending on size of plant and insect infestation.

Do not apply to tomatoes within 10 days of harvest and to peppers within 25 days of harvest.

CAUTIONS

Do not use on forage crops to be fed to dairy animals or to animals being finished for slaughter.

Toxic to fish and wildlife. Do not contaminate any body of water nor apply to any area not specified on the label.

Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Manufactured by

Reg. U. S. Pat. Off.

FOR USE ON TOBACCO, TOMATO AND PEPPER INSECTS

ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methylcarbamate) 5%

Parathion (O,O-diethyl O-p-nitrophenyl thiophosphate) 1%

INERT INGREDIENTS

94%

100%

DANGER: Keep Out of Reach of Children.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

DANGER

Poisonous by skin contact, inhalation, or swallowing. Rapidly absorbed through the skin. Repeated exposure may, without symptoms, be increasingly hazardous. Keep all persons out of operating areas or vicinity where there may be danger of poisonous drift. Vacated areas should not be re-entered until Parathion residues have dissipated.

DO NOT GET ON SKIN OR CLOTHING OR IN EYES.

Wear natural rubber gloves and goggles and clean waterproof or freshly laundered protective clothing (coveralls, cap, etc.). Destroy and replace gloves frequently. Wash thoroughly with soap and warm water before smoking or eating. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and water before re-use. In case of contact, immediately remove contaminated clothing and wash thoroughly with soap and warm water.

DO NOT BREATHE DUST, VAPOR OR SPRAY MIST.

Wear a mask or respirator of type passed by U. S. Department of Agriculture for Parathion protection. Airplane pilots should wear full face canister type mask.

If handled indoors provide mechanical exhaust ventilation.

DO NOT CONTAMINATE FOOD OR FEED PRODUCTS. Bury spillage. Clean up area with strong lye solution. Burn or bury containers. Never re-use. Stay out of smoke or fumes.

PHYSICIANS NOTE

Warning symptoms include weakness, headache, tightness in chest, blurring vision, non-reactive pin point pupils, salivating, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT: Give atropine, preferably by injection, Grain 1/100, two or three tablets at once, and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may without warning cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.



POISON

2-PAM is also antidotal and may be administered in conjunction with atropine.

Atropine is an antidote. Consult physician for emergency supply.

ANTIDOTE

Call a physician at once in all cases of suspected Parathion poisoning. If symptoms or signs of poisoning include blurred vision, abdominal cramps and tightness of chest don't wait for a doctor but give two atropine tablets (each 1-100 gr.) at once. Remove patient immediately from area where Parathion is present. Remove contaminated clothing and wash skin clean with plenty of soap and water, to remove all traces of Parathion. If swallowed, induce vomiting by giving warm salty or soapy water and repeat until vomit fluid is clear. Do not give liquids to unconscious patient. Have patient lie down and keep quiet.

Notice of Warranty

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the seller, and buyer assumes the risk of any such use.

TRIANGLE CHEMICAL COMPANY

Macon, Georgia

USDA Reg. No. 1842-209

Net Contents _____ Gals.

FOR USE OR STORAGE IN OR AROUND THE HOME.

DANGER

us by skin contact. irrita-
allowing. Rapidly absorbed
he skin. Repeated exposure
ut symptoms, be increasing-
us. Keep all persons out of
areas or vicinity where there
nger of poisonous drift. Va-
s should not be re-entered
athion residues have dis-

DO NOT GET ON SKIN OR ING OR IN EYES.

tural rubber gloves and goggles
clean waterproof or freshly
protective clothing (cover-
etc.). Destroy and replace
quently. Wash thoroughly
and warm water before
r eating. Bathe immediately
and change all clothing.
hing thoroughly with soap
before re-use. In case of con-
diately remove contaminated
and wash thoroughly with
warm water.

DO NOT BREATHE DUST, VAPOR OR SPRAY MIST.

ANGLE CHEMICAL COMPANY

Net Contents _____

Gals.

Wear a mask or respirator of type
passed by U. S. Department of Agri-
culture for Parathion protection. Air-
plane pilots should wear full face can-
ister type mask.

If handled indoors provide mechan-
ical exhaust ventilation.

**DO NOT CONTAMINATE FOOD
OR FEED PRODUCTS.** Bury spil-
lage. Clean up area with strong lye
solution. Burn or bury containers. Nev-
er re-use. Stay out of smoke or fumes.

PHYSICIANS NOTE

*Warning symptoms include weakness, head-
ache, tightness in chest, blurring vision, non-
reactive pin point pupils, salivating, sweating,
nausea, vomiting, diarrhea and abdominal
cramps.*

*TREATMENT: Give atropine, preferably
by injection, Grain 1/100, two or three tab-
lets at once, and parenterally or orally every
hour as required up to 30 tablets or until
pupils dilate. Never give morphine. Clear
chest by postural drainage. Artificial respira-
tion or oxygen administration may be neces-
sary. Observe patient continuously 48 hours.
Repeated exposure to cholinesterase inhibi-
tors may without warning cause prolonged
susceptibility to very small doses of any
cholinesterase inhibitor. Allow no further ex-
posure until time for cholinesterase regenera-
tion has been allowed as determined by blood
tests.*

Z-FAM is also antidotal and may be
administered in conjunction with
atropine.

Atropine is an antidote. Consult
physician for emergency supply.

ANTIDOTE

Call a physician at once in all cases
of suspected Parathion poisoning. If
symptoms or signs of poisoning include
blurred vision, abdominal cramps and
tightness of chest don't wait for a
doctor but give two atropine tablets
(each 1/100 gr.) at once. Remove
patient immediately from area where
Parathion is present. Remove contami-
nated clothing and wash skin clean
with plenty of soap and water, to re-
move all traces of Parathion. If swal-
lowed, induce vomiting by giving warm
salty or soapy water and repeat until
vomit fluid is clear. Do not give
liquids to unconscious patient. Have
patient lie down and keep quiet.

Notice of Warranty

Seller warrants that this product conforms to its
chemical description and is reasonably fit for the
purpose stated on the label when used in accordance
with directions under normal conditions of use, but
neither this warranty nor any other warranty of
merchantability or fitness for a particular purpose,
expressed or implied, extends to the use of this
product contrary to label instructions, or under
abnormal conditions, or under conditions not rea-
sonably foreseeable to the seller, and buyer assumes
the risk of any such use.

Macon, Georgia

USDA Reg. No. 1842-209