# Wettable Sulphur

For Ornamentals, Flowers, and Vegetables

## **ACTIVE INGREDIENTS:**

Sulphur as elemental INERT INGREDIENTS

94.0%

6.0%

100.0%

CAUTION: Keep out of reach of children

Net Contents 4 Lbs.



# WETTABLE SULPHUR

### Directions for Use

IRIANGLE WEITTABLE SULPHUR may be used as a dust or spray. In most cases, dusting will prove satisfactory. If used as a spray, add six level tablespoons per gallon of water. Apply to thoroughly cover, at 5 to 7 day intervals beginning when the disease first appears. Continue applications until all danger of disease is past.

During rainy weather it may be necessary to apply every 2 to 3 days. Use as above for:

POWDERY MILDEW on Roses, Dogwood, Perennials, Annuals, and Ornamentals.

RED SPIDER on Arborvitae, Cedar, Hydrangea, Rose, Beans, Strawberry, and Turnips.

 $\sim 2.5 \times VN/ROT$  and SCAB on Peaches and Plums.

 $^{++}$  , the above fruit and vegetables apply at a rate of  $\pm_2$  lb, actual  $\sin_4$  , per 1000 sq. ft.

### Caution

- I. Do not apply to cucumbers, melons, or squash,
- 2. Do not apply this material during very hot periods of the day.
- 3. Avoid contamination of feed and food stuffs.

# Notice of Warranty

Seller warrants that this product conforms to its cherrical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use but neither this warranty nor any other warrants of merchantability or latiness for a particular purpose expressed or implied extends to the use of this product contrary to laber instructions, or under abnormal conditions or under conditions not reasonably fore secrole to the seller and buyer issuities the risk of any such use

USDA Reg. No. 1842-212

