

30% NEMAGON GRANULES

ACTIVE INGREDIENTS:

1,2-Dibromo-3-Chloropropane

30.0%

Other halogenated C3 Compounds

1.0%

INERT INGREDIENTS

69.0%

100.0%

WARNING: Keep Out of Reach of Children

See Additional Warnings below.

GENERAL DIRECTIONS: TRIANGLE Nemagon Granules should be applied when soil temperatures range from 50° to 80° F at a depth of 6 inches. The soil should be in good tilth and contain adequate moisture for seed germination.

Extremely wet or dry soils should not be treated. The nemagon gradually disperses, killing Nematodes as it moves. Because of its slowness of movement, 2 or 3 months may be required to penetrate and kill Nematodes.

Directions For Use

For control of Nematodes in lawns and turf—apply ½ pound to each 100 square feet. Apply at any time with lawn fertilizer spreader or by hand, protected with rubber gloves. Immediately after application, wash thoroughly into soil with hose or sprinkler system applying at least 60 gallons of water per 100 square feet (equivalent to 1 inch of rain).

For established shrubs and small fruits such as Azalea, Camellia, Boxwood, Gardenia, Hibiscus, and Grape—apply ¼ lb. per 100 square feet and water thoroughly into soil.

Apply with fertilizer spreader or by hand protected with rubber gloves. Apply at least 60 gallons of water per 100 sq. ft. (equivalent to 1 inch of rain).

For Snap Beans, Cucumbers, Summer Squash, Carrots, Celery, Parsnips, Radishes and Turnips—

—Preplanting row application (3 ft. rows) apply in trench rows before planting at a rate of 2½ oz. Granules to 60 ft. of row (equivalent to 13 lbs. Technical Nemagon per acre). Cover trench with soil and plant seed.* For transplants, wait 2 weeks before planting. Do not feed vines of beans to livestock. Do not feed treated plant parts of carrots and parsnips to livestock.

For Cabbage, Cauliflower, Broccoli, Brussels Sprouts and Okra—apply at least at a rate of 1½ lbs. Granules to 1,000 square feet of surface (equivalent to approximately 20 lbs. Technical Nemagon per acre). Work treatment into top 6 to 8 inches of soil and plant seed.* For transplants, wait 2 weeks before planting.

For Lettuce and Escalope (Escarole)—apply evenly before planting at 6 inch depth in rows 12 inches apart at the rate of 1 lb. Granules to 200 ft. of row. This is equivalent to 16½ lbs. of actual Nemagon per acre. Fill trenches and wait 2 weeks before planting.

Honeydew Melons and Muskmelons—Preplanting treatment 6 ft. rows. Prepare a trench 6" deep and apply ¼ lb. granules to 100 ft. of row. Cover trench and plant seed.*

*Before planting seed, use a waiting period of 7 to 14 days for shallow rooted crops—4 to 6 weeks for deep rooted plants.

Cautions

READ THE DIRECTIONS ON THE LABEL AND FOLLOW THEM AT ALL TIMES.

Do not use Nemagon to treat soil to be planted to onions, garlic, celeriac, white potatoes, bell peppers, tomatoes, chrysanthemums or dwarf palms. Do not feed treated vines of beans to livestock. Confined areas such as green houses should be well ventilated during and after application.

WARNING: HARMFUL IF INHALED. MAY BURN SKIN. CAUSES IRRITATION OF EYES, NOSE AND THROAT. MAY BE ABSORBED THROUGH SKIN.

HARMFUL IF SWALLOWED OR INHALED

Do not breathe vapor. Use in well ventilated area. Do not get in eyes, avoid contact with skin or clothing. Do not take internally. Wash thoroughly with soap and water after handling and before eating or smoking. In case of accidental spillage on person or clothing, immediately remove contaminated shoes and clothing and wash skin with soap and water; flush eyes with plenty of water for at least 15 minutes and get medical attention. Wash clothing and air shoes thoroughly before re-use. Do not contaminate feed or foodstuffs. Keep out of reach of children. Keep children and pets off treated areas until this material has been washed into the soil and the treated areas have dried completely.

This product is toxic to wildlife. Granules exposed on soil surface may be hazardous to birds and other wildlife. Do not reuse container. Destroy when empty.

Manufactured by

TRIANGLE CHEMICAL COMPANY

Macon, Georgia

USDA Reg. No. 1842-218

RECEIVED JUL 11 1960