

TRIANGLE EPN-5EG

MAR 2 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 1842-264 SUBJECT
TO ATTACHED COMMENTS.

ACTIVE INGREDIENTS:

O-Ethyl O-p-Nitrophenyl phenylphosphonothioate... 55.0%
Xylene..... 32.1%

INERT INGREDIENTS 12.9%

TOTAL 100.0%



POISON



DANGER: KEEP OUT OF REACH OF CHILDREN.

Poisonous by skin contact, inhalation or swallowing. Rapidly absorbed through skin. Repeated exposure may, without symptoms, be increasingly hazardous. Do not get in eyes, on skin, or clothing. Do not breathe vapor or spray mist. Wear clean, heavy natural rubber gloves, rubber boots, and clean waterproof or freshly laundered protective clothing (coveralls, cap), and goggles when applying and handling. Destroy and replace gloves frequently. Wear an effective mask or respirator of a type passed by the U. S. Bureau of Mines for EPN protection. Do not contaminate feed or foodstuffs. Wash hands, arms, and face thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and warm water; for eyes, flush with water for 15 minutes and get medical attention. Keep this product away from heat and open flame. **PROTECT WILDLIFE:** This product is poisonous to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply in any manner not specified on the label.

ANTIDOTE AND FIRST AID TREATMENT

Atropine is an antidote. Consult physician for emergency supply. Call a physician at once in all cases of suspected poisoning. Never give anything by mouth to an unconscious person. **INTERNAL:** Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **EXTERNAL:** Immediately remove all contaminated clothing and wash skin thoroughly with soap and warm water.

NOTE TO PHYSICIAN

Repeated therapeutic doses of Atropine may be effective. 2-PAM is also antidotal and may be administered in conjunction with Atropine. Do not use morphine. **SYMPTOMS OF POISONING** are digestive upsets, nausea, vomiting, cramps, feeling of pressure under breast bone, diarrhea, sweating, excessive salivation, restlessness, riddiness, headache, muscle tremors, twitching

TREATMENT: Remove person from area of poisoning. Give two 1/100 grain atropine tablets. Do not resume operations with any organophosphorous insecticide, including EPN, until the physician has tested a sample of your blood for organophosphorous effect and tells you it is safe to do so.

DIRECTIONS

WARNING: This product should be used only by experienced or trained operators properly protected from the hazards involved. Use in concentrate orchard sprayers increases the hazard. Do not apply or allow to drift to areas occupied by unprotected humans or animals. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. If this product is spilled, wash down with large amounts of water. Decontaminate areas and equipment by washing with a concentrated solution of alkali and detergent and rinse with water.

DECONTAMINATE AND DESTROY CONTAINER: Rinse the empty container with cold water and pour rinsings into spray tank. Crush and bury empty container. Do not contaminate water by cleaning of equipment or disposal of waste. Do not use, pour, spill or store near heat or open flame. This product is toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Extension Service.

PEACHES: To control Plum Curculio and Oriental Fruit Moth use 1 to 1 1/2 quarts in 500 gallons of water and apply at the rate of 150 to 200 gallons per acre. In concentrate sprayers or by airplane apply at the rate of 1/3 to 1/2 quart per acre diluted with sufficient water for coverage. To control Peach Tree Borer and Lesser Peach Tree Borer use 1 1/4 to 2 1/2 quarts in 500 gallons of water and apply to thoroughly wet the scaffold limbs and trunk; usually 50 gallons of spray per acre will be required. Treatments for Oriental Fruit Moth and plum curculio should be made as directed by state spray schedules, usually beginning with the first cover spray following bloom and repeated as

necessary for control. Borer sprays should be applied as directed by your state spray schedule, usually after harvest and at 30 day intervals for 2 to 4 sprays. Do not apply within 21 days of harvest.

PECANS: To control Hickory Shuckworm, use 2 quarts in 500 gallons of water and apply at the rate of 250 to 500 gallons per acre. For airplane or concentrate sprayers apply at the rate of 1 to 2 quarts per acre diluted with sufficient water for coverage. Make three late applications beginning about shell hardening time (usually about August 15). Apply second and third spray at two week intervals. To control Pecan Weevil, use at the same rate as for Hickory Shuckworm. Applications directed against the Hickory Shuckworm will also control Pecan Weevil. Where the Pecan Weevil is the only pest to be controlled only two applications are necessary. The first should be made when a minimum of six weevils can be jarred from the branches of any one tree. This is usually about August 1 to 10. The second application should be made 10 days to two weeks later. To control Aphids, Mites, and Pecan Nut Casebearer use 1 1/2 quarts in 500 gallons of water or 1 to 1 1/2 quarts per acre in sufficient water for coverage if applied by airplane or concentrate sprayer. To control nut casebearer, make application at the time small nuts have turned brown. To control Aphids and Mites, make application as soon as observed or when injury is first seen. To control Leaf Casebearer, use 2 quarts in 500 gallons of water applied at the rate of 250 to 300 gallons per acre or 1 to 1 1/4 quarts per acre in sufficient water for coverage by airplane or concentrate sprayer. Begin applications when insects or injury first appears and repeat as necessary for control. Do not apply within 21 days of harvest. Do not graze livestock in treated groves.

NOTICE

DISCLAIMER OF WARRANTY, MERCHANTABILITY AND LIABILITY - If for any reason this product is not used, applied, stored, transported or disposed of in accordance with the instructions set forth on this label, then and thereafter, Triangle Chemical Co. terminates, ends and excludes all warranties, expressed or implied, including the warranty of merchantability and fitness.

NET _____ GALS.

EPA Reg. No. 1842-264

Manufactured by **TRIANGLE CHEMICAL COMPANY** Macon, Georgia