

DIRECTIONS FOR USE

FOR CONTROL OF WIREWORMS IN TOBACCO: Use at the rate of 10 ounces to each 50 gallons of setting water.

VEGETABLE CROP PESTS: PLANTERS 40% WETTABLE CHLORDANE is suggested for use against the following vegetable pests: Harlequin Bugs on Cabbage and Collards — Do not apply after heads or other edible parts begin to form — Onion Thrips, Colorado Potato Beetles on Potatoes and Tomatoes, Horn Worms and Flea Beetles on Tomatoes. Use at the rate of 2½ pounds in 100 gallons of water at 10 to 14 day intervals with as many applications as are needed to control these insects. For Cabbage Worms, use at the rate of 5 pounds per 100 gallons of water. Apply when the eggs hatch. For Cabbage Maggots, use at the rate of 5 pounds per 100 gallons, applying to the lower parts of the stems and the ground around the plants. Do not apply to Cabbage after the heads or other edible parts begin to form. On tomatoes, do not apply after fruit begins to form.

STRAWBERRIES: For the control of Spittle Bugs, use at the rate of 2½ pounds per 100 gallons of water two weeks before blossoming. Do not apply after fruit begins to form.

PEACHES: To control Plum Curculio and Catfacing insects, use at the rate of 2½ pounds per 100 gallons of water, applying in shuck split and cover sprays as recommended by your State Agricultural Experiment Station. Do not apply within 30 days of harvest.

SOIL PESTS: For ants infesting golf courses and lawns, use at the rate of 5 ounces in 75 to 100 gallons of water and apply to 1,000 sq. of turf. For other soil and lawn pests, such as Chinch Bugs, Japanese Beetle Grubs, Mole Crickets, White Fringe Beetle Larvae, White Grubs, and Wireworms, use at the rate of 25 pounds per acre as a spray using enough water to soak the soil.

(Continued on Right Panel)

PLANTERS

40%

ACCEPTED

JUL 9, 1970

UNDER THE FEDERAL INSECTICIDE ACT
FUNGICIDE AND ROENTGEN
FOR ECONOMIC POISON
ED UNDER NO. 100-100000
TO ATTACHED COMMENTS

WETTABLE

CHLORDANE

ACTIVE INGREDIENTS:

Technical Chlordane*

10%

INERT INGREDIENTS

60%

TOTAL

100%

*Equivalent to 21% Octachloro-1,7-methanotetrahydroindane and 16% related compounds.

CAUTION KEEP OUT OF REACH OF CHILDREN.

USDA REG. NO. 1868-41



IN BUILDING: Apply as a residual spray in barns, stables, kennels, hog pens, and similar places where water staining and visible residue are not objectionable. It is effective for controlling Flies, Mosquitoes, Brown Dog Ticks, Fleas, Cockroaches and Bed Bugs. Use at the rate of 10 pounds to 25 gallons of water and apply to areas frequented by insects and in their runways, giving particular attention to cracks and crevices where they hide and also to sleeping quarters of animals, especially the cracks in nests and stalls. Repeat applications as often as necessary.

NOTE: Do not apply to plants that are to be used directly as food less than 30 days prior to harvest. Do not use on dairy cattle or inside of dairy barns. Do not apply in poultry houses. Do not use on forage crops to be fed dairy animals or those being finished for slaughter.

CAUTION

Harmful if swallowed. Contact with skin causes toxic symptoms. Avoid contamination of feed and food-stuffs. Avoid breathing spray mist.

Keep children and pets off treated areas until this material is washed into soil and grass is dry. In case of contact with skin, wash with soap and water. This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Birds feeding on treated areas may be killed. Apply this product as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse container. Destroy when empty.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

DIRECTIONS FOR USE

TO CONTROL WIREWORMS IN IRISH OR WHITE POTATOES AND SWEET POTATOES: Preplant only
broadcast soil application, apply with any standard broom sprayer in sufficient water to give uniform coverage, making sure to disk or otherwise mix well into top several inches of soil. The dosages rates are as follows: On sandy soils use 2½ pints of this product per acre; on clay or loams use 1½ quarts per acre; on peats or musks use 2 to 3 quarts per acre.

Planters 75% Chlordane Emulsifiable Concentrate will also control Mole Crickets, Cutworms, Seed Corn Maggots, White Grubs, Tuber Flea Beetle Larvae, European Chafer Larvae, White Fringed Beetle Larvae and Earwigs. The maximum dosage for any preplant broadcast soil application is 5 quarts (10 lbs. actual Chlordane) per acre.

WARNING

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs.

Do not use in commercial food preparation areas or in edible products areas of food processing plants.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

HOUSEHOLD APPLICATIONS — PLANTERS 75% CHLORDANE EMULSIFIABLE CONCENTRATE can be diluted as directed below and sprayed as a coarse spray or brushed on cracks or other areas such as window sills, floor molding, water pipes, and other natural habitats and hiding places of the insect pests. In interior situations, this product will give residual control to the treated premises against re-infestation. Surfaces exposed to weathering or washings should be treated periodically as needed.

Add 1½ tablespoons of PLANTERS 75% CHLORDANE EMULSIFIABLE to one quart of water, stir, and apply as follows:

TO CONTROL ROACHES OR WATERBUGS: Thoroughly treat infested cracks and other hiding places. Apply locally on exposed surfaces where roaches will crawl when they come out of hiding. Repeat treatment when necessary.

TO CONTROL ANTS: Thoroughly treat ant trails and areas around door sills and window frames where the pests may enter premises. It is frequently desirable to treat the openings around water pipes, heat ducts, electrical outlets and baseboards where ants may come out into rooms from wall spaces and partitions. Repeat treatments as often as necessary to maintain effective control.

TO CONTROL SPIDERS: Apply as wet spray along infested baseboards and corners. In basements, treat pipes, storage localities, and other areas infested. Repeat treatment where necessary.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

USDA REG. NO. 1868-178

PLANTERS

75% CHLORDANE EMULSIFIABLE CONCENTRATE

CONTAINS 8 LBS. TECHNICAL
CHLORDANE PER GALLON

ACTIVE INGREDIENTS:

Technical Chlordane*	75.00%
Aliphatic Petroleum Hydrocarbons	20.00%

INERT INGREDIENTS:	5.00%
--------------------	-------

TOTAL	100.00%
-------	---------

*Equivalent to 45% Octocloro — 4,7-methano-tetrahydroindane and 30% related compounds.

WARNING KEEP OUT OF REACH OF CHILDREN

SEE OTHER WARNINGS ON SIDE PANEL

T-H