

## DIRECTIONS

**PEACHES:** For control of plum curculio, oriental fruit moth and aphids use 1 pint to each 100 gallons of water. Two to six applications are usually necessary for curculio.

**APPLES:** For control of plum curculio use 1½ pints in 100 gallons of water. For Red Spider mites use 1 pint in 100 gallons of water; two applications at 10 day intervals may be necessary. For Red Banded Leaf Roller use 1½ pints in 100 gallons of water. Make first application in first cover spray. If second brood appears, application should be made about the time of the fourth cover spray.

**TOBACCO:** For aphids, midge larvae, slugs, snails, vegetable weevil larvae, and green June Beetle larvae in plant beds use 1 pint in 100 gallons of water and apply at the rate of 3 to 10 gallons to each 100 square yards of bed. If applications are made 5 to 7 days prior to pulling plants, handlers should wear gloves. For aphids and suck flies on field plants use 1 pint in 60 gallons of water and apply at the rate of 40 to 80 gallons per acre. Applications should not be made within five days of priming or 15 days of cutting. Avoid plant juices coming in contact with the skin or other parts of the body of those who are engaged in cutting the crop.

**ORNAMENTALS (Outdoors):** For aphids, mites, mealybugs, leaf tiers, white flies, and lace bugs use 1 pint in 100 gallons of water. Do not contaminate ornamental fish ponds.

**NOTE:** Do not apply to fruits closer than 14 days before harvest.

Planters 2 Lb. Parathion Emulsion is compatible with DDT, Sulphur, neutral coppers, and many other commonly used insecticides and fungicides which are non-alkaline in nature. Consult Federal or State Experimental Stations for further information as to the compatibility of other insecticides or fungicides. Parathion also controls a large number of other insects on many crops. Get in touch with Planters Chemical Corporation or consult local authorities as to rates and proper timing.

## CAUTION

**INFLAMMABLE.** Keep away from fire. This product may injure certain varieties of apples such as McIntosh and related varieties. Do not apply to tobacco within 15 days of harvest.

# PLANTERS 2 LB. PARATHION EMULSION

Contains 2 Lbs. Technical Ethyl Parathion Per Gallon

**POISON**



### ACTIVE INGREDIENTS:

Parathion*	23.00%
Xylene	67.00%
INERT INGREDIENTS:	10.00%
<b>TOTAL</b>	<b>100.00%</b>

\*O,O-diethyl O-p-nitrophenyl thiophosphate

**DANGER** KEEP OUT OF REACH OF CHILDREN.

USDA REG. No. 1868-206

See side panels for Antidote, Caution and Warning statements.

NORFOLK, VA.



ALBANY, GA.

FAYETTEVILLE, N. C.

NET CONTENTS — GALLONS