

FARMCRAFT[®]

PERTHANE 4E

Active Ingredients:

PERTHANE Diisopropylbenzene oxide
RELATED REACTION PRODUCTS

AROMATIC PETROLEUM SOLVENTS

Inert Ingredients:

CONTENTS: 4% PERTHANE 4E

EPA Reg. No. 1871-73

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contact with skin. If skin contact occurs, wash thoroughly with soap and water after rinsing.

USE PRECAUTIONS

This product is for professional use only. Keep out of reach of children. Do not contaminate food or feed.

This product is to be applied by a registered applicator. Do not apply directly to active ingredients. Always read and follow the directions on the label.


Do not reuse empty containers. Do not drink from empty containers. Do not use by perforating a hole in the bottom of the container.

DO NOT USE ON FOOD CROPS OR VEGETABLES.
NEVER USE ON HUMAN BEINGS.

REGULATIONS FOR USE IN THE NORTWEST AND OTHER AREAS
DANGER: Do not spray on or near people, pets, or animals. Do not spray on sprayer tank and portable sprayers. Continue spraying until spray is dry. Do not apply to food crops. Do not allow to stand, reconstitute, mix or dilute before use. Use from 10 to 100 gallons of water per acre in ground sprayer and 5 to 10 gallons in aerial. Consult the State Agricultural Extension Service for additional usage information.

Broccoli, Cabbage, Cauliflower, Brussels Sprouts, to control Cabbage loopers, Imported Cabbage worms apply 1 quart per acre in 20 to 100 gallons of water. Apply when insects first appear and repeat as necessary to maintain control. Do not apply within 3 days of harvest.

Cherry, peach, plum, apricot, fruit fly apply 1 quart in 100 gallons of water. Apply when larvae first appear and repeat as necessary to control adults. Do not apply within 2 days of harvest.



Manufactured by
FARMCRAFT, INC.,
Medford, Oregon 97223

