BAG FACSIMILE



PRINTED LABEL ACCEPTED

GLUED

JUN 2 9 1973

UNDER THE FEDERAL ACTE
FUNGICIDE AND REDIM FOR ECONOMIC PORCH
ED UNIVER NO.
TO BAG

HERE

/// 1787

### 50 LBS. NET WT.

Product Farmer of the book of the experience and of storage, handling or use. It take no was the extensions of a mighod, with respect to may effect of the fit of its north and the for fitness for any purpose you specifically directed by the deli-

It, the noter passed of the taken passession, the R ver couple and accumulate respect dusty for taken, having and no of this produce in and manner a to demand all terrors and these from him or dusting not agreed both having and about to large and the course of the product of

FARMCRAFT, INC.

GRESHAM 97030 TIGARD 97223 HILLSBORO 97123 OREGON

TO ATTACHED COMMENT

HERE

## 50 LBS. NET WT.

Because Farnicraft. Inc. has no control over conditions of storage, handling or use, it makes no warranty either expressed or implied, with respect to any effect of this product, its merchantability and/or fitness for any purpose not specifically directed by its label.

By the act of purchase ard/or taking possession, the Bover accepts and assumes all responsibility for storage, handling and use of this product in such manner as to safeguard all persons and things from harm or damage, and agrees to hold harmless and ordennify Farmeraff, Inc. or other purveyor from any judyment, costs, expenses and/or Attorney's few memory in whole or part from practices by the purchaser or his transferce not in strict accordance with directions given herewith.

# FARMCRAFT, INC.

GRESHAM 97030 TIGARD 97223 HILLSBORO 97123 OREGON Cameraman's Note

# CAPTAN 7.5/ THIOD AN 3

EPA Reg. No. 1871-80

SUGGESTIONS FOR USE IN THE PACIFIC NORTHWEST: For combination plant disease and insect control as listed below apply uniformly through ground or aircraft duster before disease infection occurs, or when insects first appear. Repeat as necessary (within time limitations) to maintain control.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STRAWBERRIES, to control Botrytis rots, Omnivorous leaf tier (Strawberry fruitworms), Cyclamen mites and Spittlebugs, apply 40 to 50 pounds per acre. Do not apply within 4 days of harvest. Do not reapply within 35 days when fruit is present.

#### Active Ingredients:

### CAUTION

#### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust or contact with skin, eyes or clothing. In case of contact, immediately remove contaminated clothing and wash with soap and water. Flush eyes repeatedly with water and get medical attention. Wash throughly with soap and water after exposure, especially before smoking or eating. Launder all protective clothing before re-use. Avoid contamination of feed or foodstuffs. Do not allow this product to contaminate and adjacent crop, or areas occupied by unprotected humans or useful animals.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.