

FACSIMILE

FARMCRAFT[®]

DUST


ATTACHED

PRINTED LABEL

GLUED

TO BAG

HERE



50 LBS. NET WT.

FARMCRAFT, INC.

GRESHAM 97030 TIGARD 97223 HILLSBORO 97123
OREGON

Camera man's note
Poor Copy

TO BAG

HERE

JUN 29 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. _____ SUBJECT
TO ATTACHED COMMENTS.

1871-82

50 LBS. NET WT.

Because Farmcraft, Inc. has no control over conditions of storage, handling or use, it makes no warranty, either expressed or implied, with respect to any effect of this product, its merchantability and/or fitness for any purpose not specifically directed by its label.

By the act of purchase and/or taking possession, the Buyer accepts and assumes all responsibility for storage, handling and use of this product in such manner as to safeguard all persons and things from harm or damage, and agrees to hold harmless and indemnify Farmcraft, Inc. or other purveyor from any judgment, cost, expense and/or Attorney's fees incurred in whole or part from practices by the purchaser or his transferee not in strict accordance with directions given herewith.

FARMCRAFT, INC.

GRESHAM 97030 TIGARD 97223 HILLSBORO 97123
OREGON

TOXAPHENE 10

EPA Reg. No.
1871-82

DIRECTIONS FOR USE IN THE PACIFIC NORTHWEST: apply uniformly through ground or aircraft duster when insects first appear. Repeat as necessary (within time limitations) to maintain control.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

ALFALFA (Seed Crop), to control Alfalfa caterpillar, Lygus bug, Spittle bug, Thrips, Grasshoppers, apply 30 lbs. per acre. No time limitation.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, to control Imported Cabbageworm, Cutworms, Cabbage Looper, Diamondback moth, apply 40 to 60 lbs. per acre. Do not apply after edible parts begin to form.

BEANS (Green & Snap), to control Lygus bug, Thrips. Climbing cutworms apply 40 lbs. per acre. Do not apply after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

CORN, to control Climbing cutworms and Thrips apply 40 lbs. per acre. Do not feed treated forage to dairy animals or animals being finished for slaughter.

Active Ingredient:

Toxaphene [technical chlorinated camphene
(chlorine content 67% - 69%)] . . 10.0 %

Inert Ingredients: 90.0 %

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin, eyes or clothing. May be absorbed through skin. Wash thoroughly with soap and water after exposure, and before smoking or eating. Avoid contamination of feed or foodstuffs.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.