

BAG FACSIMILE

FARMCRAFT[®] DUST

ATTACHED

PRINTED LABEL

GLUED

TO BAG

HERE

ACCEPTED

JUL 2 1973

1871-53

50 LBS. NET WT.

FARMCRAFT DUST is a registered trademark of Farmcraft, Inc. © 1973 Farmcraft, Inc. All rights reserved. This product is made in the U.S.A. from the finest quality ingredients. It is a complete and balanced feed supplement for all types of livestock. It contains all the essential vitamins and minerals for the health and well-being of your animals. It is also a natural source of energy and helps to improve the overall condition of your stock. For more information, contact your local Farmcraft distributor or write to Farmcraft, Inc., P.O. Box 100, Hillsboro, Oregon 97123.

FARMCRAFT, INC.

GRESHAM 97030 TIGARD 97223 HILLSBORO 97123
OREGON

Camera man's note
Poor Copy

1871-83

50 LBS. NET WT.

Because Farmcraft, Inc. has no control over conditions of storage, handling or use, it makes no warranty, either expressed or implied, with respect to any effect of this product, its merchantability and/or fitness for any purpose not specifically directed by its label.

By the act of purchase and/or taking possession, the Buyer accepts and assumes all responsibility for storage, handling and use of this product in such manner as to safeguard all persons and things from harm or damage, and agrees to hold harmless and indemnify Farmcraft, Inc. or other purveyor from any judgment, cost, expenses and/or Attorney's fees incurred in whole or part from practices by the purchaser or his transferee not in strict accordance with directions given herewith.

FARMCRAFT, INC.

**GRESHAM 97030 TIGARD 97223 HILLSBORO 97123
OREGON**

SEVIN 5

FP. REG. NO.
1871-83

SUGGESTIONS FOR USE IN THE PACIFIC NORTHWEST: Apply uniformly through ground or aircraft duster when insects first appear. Repeat as necessary (within time limitations) to maintain control.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning equipment or disposal of wastes. Apply this product only as specified on this label.

BEANS, to control Beet Armyworm and Western spotted cucumber beetle, apply 40 lbs. per acre. No time limitation.

CORN, to control Corn Earworm apply 30 to 40 lbs. per acre as soon as silks appear and repeat every 3 days during pollination. For insects attacking silks and ears apply at 2 to 3 day intervals starting when silks first appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestations. Application of carbaryl to the tassel region of corn during the pollen shedding period will seriously reduce bee population.

POTATOES, to control Fleabeetle apply 40 lbs. per acre. Foliage application only.

RASPBERRIES, BLACKBERRIES, to control Leafrollers, Leafhoppers, and Orange Tortrix, apply 40 lbs. per acre. Do not apply within 7 days of harvest.

STRAWBERRIES, to control Leafrollers and Spittlebug apply 40 lbs. per acre. Do not apply within 1 day of harvest.

Active Ingredient:

Carbaryl (1-naphthyl N-methylcarbamate) 5.0 %

Inert Ingredients: 95.0 %

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Avoid inhalation of dust. Avoid contact with the skin. Wash thoroughly after exposure. Avoid contamination of feed or foodstuffs.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.