

1903-19

05/10/2004

1/1

ACCEPTED

MAY 10 2004

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 1903-19

UC Mite & Lice Bird Spray# C338 03/12/2004 09:29 AM Page 2

8 in 1

Mite & Lice Bird Spray

FOR USE ON ALL BIRDS: PARAKEETS, CANARIES, AND OTHER CAGED BIRDS

ACTIVE INGREDIENTS:
Pyrethrins 0.030%
Piperonyl Butoxide, Technical* 0.300%

INERT INGREDIENTS 99.670%

*Equivalent to 0.240% (butylcarbitol) (6-propylpiperonyl) ether and 0.06% related compounds.

CAUTION:
KEEP OUT OF REACH OF CHILDREN.
See back panel for additional precautionary statements.

NET 8 FL. OZ. U.S. (237 ml)

For All Birds

1. HARMFUL IF SWALLOWED. Avoid inhalation of vapors. Avoid contact with skin. Wash contaminated skin with soap and water. Avoid spraying food and foodstuffs. Cover fish aquariums before spraying.

2. It is a violation of Federal law to use this product inconsistent with its labeling.

For control of bird lice on parakeets, canaries and other caged birds, leave birds in cage. Remove food and foodstuffs. Spray directly on birds from a distance of 12 to 18 inches. Avoid the bird's eyes and face. Spray lightly utilizing 3 or 4 bursts. For cage mite control, remove bird(s), food and water cups, etc., and spray cage thoroughly by holding unit at least one foot from surface utilizing 5 or 6 bursts. To control mites and bird lice, may repeat treatment on cage or bird no more than two times a week. Bird cages should be cleaned in hot water as part of weekly regimen.

3. Store in a cool dry area away from heat or open flame.

4. Do not reuse container. Place in trash or offer for recycling if available. Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

Serving Your Pet Product Needs For Over 100 Years!

Eight to One Pet Products

Houpage, NY 11788

www.eightonepet.com (800) 645-5154

Made and Printed in U.S.A. Item# C338 (41-6240-05)

EPA Reg. No. 1903-19 EPA Est. No. 19030701



0 26851 00338 0