

UltraCare
 For All Birds

8 in 1

Traitement en vaporisateur contre les problèmes de parasites aviaires

Mite & Lice

Bird Spray

FOR USE ON CAGED BIRDS TO KILL MITES AND LICE

ACTIVE INGREDIENTS:	
Pyrethrins	0.03%
Piperonyl Butoxide, Technical*	0.30%
INERT INGREDIENTS:	
	99.67%
	100.00%

*Equivalent to 0.240% (butylcarbityl) (6-propylpiperonyl) ether and 0.06% related compounds.

CAUTION:
KEEP OUT OF REACH OF CHILDREN
 See back panel for additional precautionary statements

UltraCare



NET 8 FL. OZ. (237 ml)

8 in 1 MITE AND LICE BIRD SPRAY

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

CAUTION: Harmful if swallowed. Avoid inhalation of vapors. Avoid contact with skin. Wash contaminated skin with soap and water. Avoid spraying food and foodstuffs. Cover fish aquariums before spraying.

It is a violation of Federal law to use this product inconsistent with its labeling.

For control of bird lice on parakeets, canaries and other caged birds, leave birds in cage. Remove food and foodstuffs. Spray directly on birds from a distance of 12 to 18 inches. Avoid the bird's eyes and face. Spray lightly utilizing 3 or 4 bursts.

For cage mite control, remove bird(s), food and water cups, etc., and spray cage thoroughly by holding unit at least one foot from surface utilizing 5 or 6 bursts. To control mites and bird lice, may repeat treatment on cage or bird no more than two times a week. Bird cages should be cleaned in hot water as part of weekly regimen.

Store in a cool dry area away from heat or open flame.

Do not reuse container. Place in trash or for recycling if available.

Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

FIRST AID

If swallowed:

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

If in eyes:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

If inhaled:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling Poison Control Center or Doctor or going for treatment. For emergency information, call (800) 645-5154

Serving Your Pet Product Needs For Over 100 Years!

Eight to One Pet Products

Hempstead, NY 11788

www.eighttoone.com (800) 645-5154

Made and Printed in U.S.A. Item# C338 (041-0340-03)

EPA Reg. No. 1985-91 EPA Est. No. 1985-91-01



ACCEPTED

JUN 29 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 1903-19

1903-19

06/29/2004

1/1