

FOR SWIMMING POOLS . . . SANITIZER
 FOR LOCKER ROOMS . . . FOR USE IN INDUSTRIAL
 RECIRCULATING WATER COOLING TOWERS . . .

SWIMMING POOL ALGAECIDE:

1. If algae growth is observed, add 1 gallon ALGAECIDE for each 10,000 gallons of water to kill and control those algae species most commonly found in swimming pools.
2. When existing growth is dead and algae have withered, clean the pool, draining and refilling if necessary. Add 1 gallon ALGAECIDE for 25,000 gallons of water.
3. If algae growth is absent, use no less than 1.5 gallons ALGAECIDE for each 50,000 gallons of water.
4. Add 2 quarts of ALGAECIDE per 50,000 gallons of water every 5-7 days to maintain between 2 and 5 parts per million of active ingredient. See service-man for test kit.

POOL SIDE SANITIZER OF HARD SURFACES:

After cleaning surfaces, use 1.5 ounces of ALGAECIDE to 4 gallons of water. With this solution, wipe, mop or scrub wet surfaces thoroughly.

EPA Reg. No. 1926-40
 EPA EST NO. 1926-MO-1



FOR SWIMMING POOLS . . . SANITIZER
 FOR LOCKER ROOMS . . . FOR USE
 IN INDUSTRIAL RECIRCULATING
 WATER COOLING TOWERS

ACTIVE INGREDIENTS: 7.26%
 n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
 dimethyl benzyl ammonium chlorides . . . 3.63%
 n-Alkyl (50% C₁₂, 30% C₁₄, 17% C₁₆, 3% C₁₈)
 dimethyl ethylbenzyl ammonium chlorides . . . 3.63%
 INERT INGREDIENTS: 92.74%

WARNING: KEEP OUT OF REACH OF CHILDREN
 See side Panel for Other Precautions

Sold by

navy brand®
 MANUFACTURING CO

INDUSTRIAL RECIRCULATING WATER COOLING TOWERS:

When ALGAECIDE is used as directed, it will kill and prevent development of *Chlorella pyrenoidosa* (green), *Phormidium inundatum* (black algae), *Phormidium retzii* (Square D algae), *Lyngba versicolor* (blue-green).

Begin treatment when the system is in jeopardy of becoming affected, or after cleaning systems whose efficiency is already impaired. For best results, rid system of organic debris if possible. If soap or anionic detergent is used, rinse thoroughly before charging with ALGAECIDE.

INITIAL DOSE: Apply 0.28 to 0.55 gal. per 1000 gal. of water in the system (20-40 ppm of active ingredient). Apply at a point in the system where the product will be uniformly mixed. Repeat until control is achieved.

PERIODIC DOSE: When control is evident, add 0.07 to 0.21 gal. per 1000 gal. of water in the system, (5-15 ppm of active ingredient) about every 7 days, or sooner, should visible growth appear. Apply at a point in system where the product will be uniformly mixed.

WARNING:
KEEP OUT OF REACH OF CHILDREN

Causes eye or skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician.

This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds or public water.

Apply this product only as specified on this label.
 Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

ACCEPTED

MAY 23 1975

UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTRATION
 UNDER NO. 1926-40

Net Contents: U.S. Gallons