

DIRECTIONS FOR USE

FOR THE SICK ROOM: All utensils, such as bed-pans and urinals, should be washed with a solution of three ounces of disinfectant to one-half gallon of water.

WATER CLOSETS and URINALS: Use three ounces of disinfectant to one-half gallon of water. Apply freely to all surfaces to be disinfected and scrub the exposed areas.

GARBAGE: To deodorize, sprinkle with solution one-half cup to a gallon of water. Wash the empty can with some of the solution and leave the wash in can.


FOR GENERAL DISINFECTING

For general disinfecting of surfaces such as walls, floors, and woodwork, remove all filth and dirt from the premises and use a solution of one part disinfectant in 100 parts of water. This mixture acts as a disinfectant against *S. typhosa*. Apply freely to all surfaces and scrub with a mop, brush, or cloth.

In the typhoid sick room, for disinfecting hard surface areas such as bed frames, bed pans, and urinals, use one-half cup of the disinfectant to two gallons of water in a mop bucket for mopping, scrubbing, or cleaning purposes.

EPA REG. NO. 1926-70
EPA EST. NO.

5/75



NAVYPINE

PINE ODOR
Disinfectant

Phenol Coefficient 5 by A.O.A.C. Method

ACTIVE INGREDIENTS		26.5%
Isopropyl Alcohol	10.70%	
Soap	5.55%	
Pine Oil	5.00%	
Ortho-phenylphenol	3.00%	
Ortho-benzyl-para-chlorophenol	2.25%	
INERT INGREDIENTS		73.5%

Warning: KEEP OUT OF REACH OF CHILDREN

See back panel for additional precautionary and First Aid Statements

Sold by

navy brand®
MANUFACTURING CO.

This disinfectant can be used many places where clean and sanitary quarters are desired. A partial list of places using this type of disinfectant follows: Hotels, Schools, Institutions, Office Buildings, Jails, Public Buildings, Railroad Terminals, Theatres, Stores, Sanitariums.

WARNING:

KEEP OUT OF REACH OF CHILDREN

Causes eye or skin irritation. Do not get in eyes, on skin, or on clothing. Harmful if swallowed.

Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing.

If swallowed, drink promptly, a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.
Rinse empty container with water and discard in a safe place.

NET CONTENTS: U.S. GAL.