For use as a disinfectant-cleaner in hospitals, nursing homes, restaurants, food processing plants, schools and public buildings. NO-BAC BACTERIACIDE should always be diluted with water before using, as the use of the concentrated disinfectant causes skin irritation.

Be sure all dirt is swept up and removed before applying disinfectant, and make sure disinfectant solution comes in contact with all contaminated areas.


To disinfect floors, cellars, basements, toilet rooms, bathrooms, lavatories, urinals, sinks, clean thoroughly: use 1 oz. of this preparation to every gallon of water; mop in regular manner. If it is impractical to mop specific places, use 3 ounces per gallon of water and spray thoroughly over entire surface.

To disinfect kitchen tables, drainboards, marble, tile, linoleum, limestone, glass, rubber, soda fountains, refrigerators, cooling rooms, refrigerator cars, rinse thoroughly with a dilution of 2 ounces of this product to a gallon of water, cleansing in regular manner, allowing contact for ten minutes.

Treated food contact surfaces must be thoroughly scrubbed with detergent and rinsed with potable water prior to reuse.

For more drastic conditions such as disinfection of exceptionally dirty urinals, garbage cans, a most thorough cleaning must be done by use of strong cleaning solutions, removing all dirt, then make sure disinfectant comes in contact with all contaminated areas; disinfect with a solution of 4 ounces of disinfectant per gallon.

For use in reducing the spread of athletes foot fungion environmental surfaces, two ounces in one gallon of water is recommended for scrubbing shower room floor, locker room benches, runways, bath mats and other surfaces commonly contacted by the bare feet.



FUNGICIDE AND RODENTICIDE ACT

FOR ECONOMIC POISON REGISTER-

ED UNDER NO. / 12 1 - 7 1

NO-BAC BACTERIACIDE

| NON-SELECTIVE | DELIGHTFUL | MINT ODOR |
|----------------------------|------------|-------------|
| ACTIVE INGREDIENTS | | 39 % |
| 1. Isopropyl Alcohol | | 25.94% |
| 2. Soap | | 5.69% |
| 3. o-Benzyl-p-Chlorophenol | | 4.90% |
| 4. Methyl Salicylate | | 1.22% |
| 5. o-Phenylphenol | | 0.75% |
| 6. Essential Oils | | |
| INERT INGREDIENTS | | 61% |

WARNING: KEEP OUT OF REACH OF CHILDREN

(See First Aid and other precautions on side panel.

Sold By

navy brand ®

FOR HOSPITAL USE

For Disinfection Against Pseudomonas Aeruginosa

Mix 240 c.c. of disinfectant with each gallon of water for prophylactic disinfection of hospital equipment, furnishings, floors, rubber goods. In cold disinfection of surgical instruments, preclean to remove all adhering blood and serous exudates prior to immersion in or tree ment with the disinfecting solution. For contaminated areas, mix 320 c.c. disinfectant with each gallon of water.

TUBERCLE BACILLI: A one percent solution (1:100) of this disinfectant is recommended to combat tubercle bacilli.

WARNING KEEP OUT OF REACH OF CHILDREN

Causes eye or skin irritation. Do not get in eyes, on skin, or on clothing.

Harmful if swallowed. Avoid inhalation of fumes.

Avoid contamination of food.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

Rinse empty container thoroughly with water and discard it.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or, if these are not available, drink large quantities of water. Avoid alcohol. Call a physician.

> EPA REG. NO. 1926 71 EPA EST. NO.

> > Net Contents: U.S. Gallons