

Navy Brand Manufacturing Company
5111 Southwest Avenue
St. Louis, MO 63110-3496

DEC 19 1988

Attention: Rudolf Oppenheim

Gentlemen:

Subject: G.G. Cleaner
EPA Registration No. 1926-72
Your Submission Dated October 22, 1987
EPA Received Date October 27, 1987

The amendment referred to above, submitted in connection with the Label Improvement Program Notice for Antimicrobial Products Used as Food Contact Surface Sanitizers, is acceptable provided that you:

1. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling.
 - a. Delete all sanitization claims from your label.
 - b. Delete the word "milk."
 - c. Revise the referral statement to read:

See side panel for additional precautionary statements and Statement of Practical Treatment.

2. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with the registration will be subject to cancellation in accordance with section 6(e) of the Federal Insecticide, Fungicide, and Rodenticide Act. Your release for shipment of the product constitutes acceptance of these conditions.

50359:I:Lee:L-15:KENCO:12/6/88:1/20/89:SG:vo:ek:rw

CONCURRENCES

SYMBOL								
SURNAME								
DATE								

A stamped copy of the label is enclosed for your records.

If you have any questions, contact Karen Leavy at (703) 557-3966.

Sincerely yours,

John H. Lee
Product Manager (31)
Antimicrobial Program Branch
Registration Division (TS-767C)

Enclosures

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

DISINFECTION—To disinfect hard inanimate surfaces (such as walls, floors, table tops), add 2 ounces G.G. CLEANER detergent/disinfectant per gallon of water. Apply solution with mop, cloth, or sponge or mechanical sprayer so as to wet thoroughly. Allow to remain wet for 10 minutes and then let air dry.

To disinfect toilet bowls, flush toilet, add 2 ounces G.G. CLEANER detergent/disinfectant directly to the bowl water. Swab the bowl completely using a scrub brush or toilet mop, making sure to get under the rim. Let stand for 10 minutes and flush.

For heavily soiled areas, a pre-cleaning step is required and recommended. Prepare a fresh solution for each use.

MILDEWSTAT—To control mold and mildew on pre-cleaned, hard non-porous surfaces (such as floors, walls, table tops) add 2 ounces G.G. CLEANER per gallon of water. Apply solution with a cloth, mop, or sponge making sure to wet all surfaces completely. Let air dry. Prepare a fresh solution for each use. Repeat application at weekly intervals or when mildew growth reappears.



NOV 27 1967
1061 27 130

G.G. CLEANER

DETERGENT/DISINFECTANT

Cleaner	Disinfectant
Detergent	Fungicide
Deodorizer	(against Pathogenic Fungi)
Mildewstat	Virucide*
(on hard inanimate surfaces)	

ACTIVE INGREDIENTS:

n-Alkyl (60% C_{14} , 30% C_{12} , 5% C_{10} , 5% C_{16})
dimethyl benzyl ammonium chlorides . . . 1.25%

n-Alkyl (68% C_{12} , 32% C_{14})
dimethyl ethylbenzyl ammonium chlorides . . . 1.25%

INERT INGREDIENTS: 97.50%

TOTAL 100.00%

EPA REG. NO. 1926-72

EPA EST. NO. 1926-MO-01

KEEP OUT OF REACH OF CHILDREN

DANGER

See side panels for additional precautionary and first aid statements.

PRECAUTIONARY STATEMENTS—Hazard to Human and Domestic Animals

DANGER: Keep Out of Reach of Children. Causes severe eye and skin damage. Do not get in eyes, skin or on clothing. Harmful if swallowed. Wear goggles or face shield and rubber gloves when handling. Avoid contamination of food. Remove and wash contaminated clothing.

STATEMENT OF PRACTICAL TREATMENT: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measure against circulatory shock, respiratory depression and convulsion may be needed.