



**CREOSOTE-OIL  
BLEND  
WOOD PRESERVATIVE  
FOR WOOD POSTS  
AND DIMENSIONAL TIMBERS**

Aids in Control of Wood Rot, Decay,  
Powder Post Beetles, Carpenter Ants

ACTIVE INGREDIENT: Coal Tar Creosote Oils.....20%  
INERT INGREDIENT .....80%

NET CONTENTS: ONE GALLON

**WARNING**

**KEEP OUT OF REACH OF CHILDREN  
HARMFUL OR FATAL IF SWALLOWED  
SEE ADDITIONAL WARNINGS ON BACK PANEL**

Made by  
**ELCO MANUFACTURING CO.**  
Pittsburgh, Pa. 15215

F.P.A. REG. NO. 1941-38

**DIRECTIONS**

Wood to be treated should be dry, well seasoned, and free of bark. Frequent inspection and re-treatment may be necessary to maintain control of fungal decay.

Application can be made by Dip, Brush or Spray. Most effective method is Dipping, least effective by Spraying.

For Dipping—soak as long as possible, minimum of 5 minutes per inch of thickness and 48 hours for fence posts, with temperature of liquid as warm as practicable. The thicker the timber, the longer it should soak. For Brush or Spray—Apply two liberal coats especially at cut ends, checks and knots.

**WARNING**

Harmful if swallowed or absorbed through skin. Causes skin irritation. Do not breathe vapor, fumes or spray mist. Do not get into eyes, on skin or on clothing. Wash thoroughly with warm water and soap. Don't smoke while using. Avoid contamination of food and feedstuffs. Keep away from children. Care should be taken in application to avoid grass and shrubbery. Do not use, pour, spill or store near heat or open flames.

Do not reuse empty small container. Wash thoroughly with water and detergent. Discard in a safe place. Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

**ANTIDOTE:** In case of swallowing administer antidote such as milk, raw egg white, gruel or corn starch paste. Follow with an emetic such as a tablespoon of mustard in a glass of warm water. Call physician.

1941-38