

ACCEPTED

DEC 7 1972

UNDER THE FEDERAL INSECTICIDE
ACT AND TOXIC SUBSTANCE ACT
FOR ECONOMIC PESTICIDE REGISTERED
ED UNDER FIFD SUBJECT
TO ATTACHED COMMENTS

DIRECTIONS FOR USE:

ROACHES, WATERBUGS, ANTS, SILVERFISH:

Use 3 to 4 ounces for each 1000 cu. ft. of space, directing the spray to contact as many insects as possible. Pay particular attention to baseboards, wall mouldings, woodwork pipe openings, breeding places, and nests. Do not remain in treated areas. Ventilate the area after treatment is completed.

FLIES, MOSQUITOES, GNATS: Use $\frac{1}{2}$ ounce per 1000 cu. ft. of space, directing the spray toward the ceiling. Do not remain in treated areas. Ventilate the area after treatment is completed.

ACCESSIBLE STAGES OF GRANARY WEEVIL, RICE WEEVIL, CONFUSED FLOUR BEETLE, CADELLE, SAW-TOOTHED GRAIN BEETLE, ANGOUMOIS GRAIN MOTH, MEDITERRANEAN FLOUR MOTH AND MEAL WORM (CAUTION: Do not spray directly on exposed food, flour, grain, or cereals.) Use 3-4 ounces per 1000 cu. ft. of space, directing the spray so as to contact the insects. Pay particular attention to cracks, crevices, and other hiding places. Repeat treatment as required. Do not remain in treated areas. Ventilate the area after treatment is completed.

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid inhalation of spray mist. Wash thoroughly after handling. Avoid contamination of feed and food stuffs. Do not store near heat or flame.

Remove pets, birds and cover fish aquariums before spraying.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

When using the product in food processing plants, "Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. Apply only when the facility is not in operation."