

DILUTION GUIDE OF CHLORDANE

| PERCENT CHLORDANE IN FINISHED FORMULATION | VOLUME OF FINISHED FORMULATION (Gallons) | AMOUNT OF CHLORDANE NECESSARY |
|---|--|-------------------------------|
| 1% | 48 | 2 quarts |
| | 96 | 1 gallon |
| 2% | 1 | 2.7 ounces |
| | 5 | 13.5 ounces |
| | 48 | 1 gallon |
| 3% | 1 | 4 ounces |
| | 5 | 20.25 ounces |

CAUTION

Harmful if swallowed. Contact with skin can cause toxic symptoms. Avoid breathing vapors or spray mist. In case of contact with skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Keep out of the reach of children.

Keep out of any body of water. Toxic to fish and wildlife. Do not use in any manner not specified on this label.

ANTIDOTE

EXTERNAL—In case of spillage on skin wash promptly with soap and water.

INTERNAL—Emetic of 1 tablespoonful mustard in tumbler of water. Call a physician.

CONTAINER DISPOSAL

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. (Keep out of smoke.)

USDA Reg. No. 1441-53

ELCO-101

EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENTS:

*Technical Chlordane 72.0%

Petroleum Distillate 21.0%

INERT INGREDIENTS 7.0%

TOTAL 100.0%

*Equivalent to 43.2% Octachloro-4, 7-methano tetrahydrodane and 28.8% related compounds

Contains 8.0% lbs. actual chlordane per gallon.

FOR USE BY PROFESSIONAL PCO'S ONLY.

Net Contents

WARNING

KEEP OUT OF REACH OF CHILDREN

See left panel for antidote and additional caution statements.

Made by

ELCO MANUFACTURING CO.

Pittsburgh, Pa. 15215

1941-53
 REGISTERED
 APR 21 1971

1. 100% Pure
 2. 100% Pure
 3. 100% Pure
 4. 100% Pure
 5. 100% Pure
 6. 100% Pure

DIRECTION GUIDE FOR CYLONDAZE

| INSECT PEST | CONCENTRATION | APPLICATION |
|--|---------------|---|
| Ants Centipedes Roaches Scorpions Silverfish Spiders Crickets Waterbugs | 2.0% Spray | Thoroughly spray areas such as baseboards, closets, storage areas, openings around water pipes, air conditioning ducts, window frames, wall cracks, electrical outlets. |
| Clothes moths | 2 1/2 % Spray | Thoroughly dampen all surfaces of clothing, blankets and other woolen cloth. Treated articles should be dried and then placed immediately in storage. Dry clean before using them as clothing or bedding. |
| Mosquitoes Wasps | 2-3% Spraying | Thoroughly spray around windows and areas frequented by mosquitoes. To prevent the entry of these insects into buildings thoroughly spray window frames, doorsills and selected areas of porches where they tend to alight and congregate. Spray the wasp nest after dark when all the insects have returned to the nest. |
| Brown dog tick | 2-3% Spray | Thoroughly spray or paint with a coarse wet spray infested areas, baseboards, window and door frames, wall cracks, sleeping quarters of household pets and localized areas of floors and floor coverings. Use fresh bedding after treatment. Do not spray pets or other animals. |

DIRECTIONS: Grubs of Japanese Beetle, Rhinoceros Beetle, Asiatic Garden Beetle, Oriental Beetle, Rose, European and Masked Chafer Beetles and Ants - Use 5 quarts per acre at turf (4 ft. or less - 1,000 sq. ft.), apply in sufficient water for thorough coverage. Make application at any time when the ground is not frozen or soggy. Spraying or foliar treatments are preferred. Immediately after application work or water into top few inches of soil. Treatment will kill a few per

| | CONCENTRATION | APPLICATION |
|---|---------------|---|
| Ants Centipedes Roaches Scorpions Silverfish Spiders Cicadas Waterbugs | 2-3% Spray | Thoroughly spray around doorsills, baseboards, closets, storage cabinets, openings around water pipes, heat and air conditioning ducts, window frames, wall cracks, electrical outlets |
| Clothes moths | 2-3% Spray | Thoroughly dampen all surfaces of clothing, blankets and other woolen cloth. Treated articles should be dried and then placed immediately in storage. Dry clean before using them as clothing or bedding. |
| Mosquitoes Wasps | 2-3% Spr. | Thoroughly spray around windows and areas frequented by mosquitoes. To prevent the entry of these insects into buildings thoroughly spray window frames, doorsills and selected areas of porches where they tend to alight and congregate. Spray the wasp nest after dark when all the insects have returned to the nest. |
| Brown dog tick | 2-3% Spray | Thoroughly spray or paint with a coarse wet spray infested areas, baseboards, window and door frames, wall cracks, sleeping quarters of household pets and localized areas of floors and floor coverings. Use fresh bedding after treatment. Do not spray pets or other animals. |

DIRECTIONS: Grubs of Japanese Beetle, Rhinoceros Beetle, Asiatic Garden Beetle, Oriental Beetle, Rose, European and Masked Chafer Beetles and Ants - use 5 quarts per acre of turf (4 ft. or per 1,000 sq. ft.), apply in sufficient water for thorough coverage. Make application at any time when the ground is not frozen or soggy. Spring or fall treatments are preferred. Immediately after application work or wash into top few inches of soil. Treatment will kill a large percentage of the grubs growing in the turf at time of application and will control ant and broods for 2 or 3 years.

1941-53
 ACCEPTED
 40 81