



R. T. VANDERBILT COMPANY, INC.

20 Winfield Street, Norwalk, CT. 06855

NET WEIGHT 100 LBS.

1 AS 1000

5-18-70

100

1965-31

Vancide* PA

U.S. PATENT NOS. 3,052,597; 3,199,990; 3,471,571

PAINT MOLD INHIBITOR

ACTIVE INGREDIENTS:

Trans-1,2-bis(n-propylsulfonyl)ethene

95%

INERT INGREDIENTS:

5%

100%

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

DIRECTIONS FOR USE: Protect paint from damage by mold by milling in Vancide PA using 0.5% to 1.0% by weight of the paint. Not recommended in emulsion paints of pH 7 or above.

DANGER: Keep out of reach of children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food. Do not reuse container. Destroy when empty.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

E.P.A. REG. NO. 1965-31
EPA EST. No. 1965-NJ-01

NON-WARRANTY OF RESULTS

Our recommendations for use are based on tests believed reliable. Since the use is beyond our control, neither we nor the manufacturer can guarantee the results if such use is not in accordance with directions. We disclaim any responsibility for damages resulting from careless or improper handling or use.

EP
5.4

NET WEIGHT 100 LBS.

Vancide* PA

PAINT MOLD INHIBITOR

FIRST AID

ACTIVE INGREDIENTS:

Trans-1,2-bis(n-propylsulfonyl)ethane

95%

INERT INGREDIENTS:

5%

100%

DIRECTIONS FOR USE:

NOTE TO PHYSICIAN

DANGER: Keep out of reach of children. Corrosive.