

# ORTHO Ferbam 95 Wettable

NET WEIGHT 30 LBS.

ACCEPTED  
10-17-68  
UNDER THE FEDERAL INSECTICIDE  
AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 1965-37

READ THE LABEL

KEEP OUT OF REACH OF CHILDREN.

**CAUTION:** May cause irritation of eyes, nose, throat and skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes and clothing. Keep away from fire or sparks. In case of contact, flush with plenty of water; for eyes, get medical attention. Do not store or transport with feed or food. Keep out of reach of domestic animals and pets. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of lakes, streams and ponds.

Active Ingredient  
Ferbam (Ferric Dimethyldithiocarbamate) ..... 95%  
Inert Ingredients ..... 5%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

All Recommendations Based on 100 Gallons of Spray.

**APPLES:** Scab— $\frac{3}{4}$  to 1½ lbs. at pre-bloom, calyx and early-cover sprays;  $\frac{1}{2}$  lb. in later cover sprays. In combination with sulfur, use  $\frac{3}{8}$  lb. plus 3 lbs. wettable sulfur. Not recommended for Golden Delicious and other varieties susceptible to russetting. Rust—Use scab recommendations through first cover and include a special bloom spray of  $\frac{3}{8}$  lb. Other Diseases (Apple Blotch, Bitter Rot, Black Rot, Blossom End Rot, Brooks Spot, Fly Speck, Frog-eye Leaf Spot, Sooty Blotch)—Use scab recommendations as long as any of the diseases appear. Do not apply to Apples later than 7 days before harvest.

**ALMONDS:** Shot-Hole—1½ to 1½ lbs. at popcorn and petal-fall.

**APRICOTS:** Brown Rot, Shot-Hole—1½ lbs. at red-bud, early-bloom and full-bloom. Green or Jacket Rot—1½ lbs. at full-bloom and petal-fall. Do not apply to Apricots later than 21 days before harvest.

**BLUEBERRIES:** Mummy Berry—2¼ lbs. at loose bud scales and 7 days later. Botrytis Blight—2¼ lbs. immediately before bloom, full bloom and 2 to 3 weeks later. Do not apply later than 40 days before harvest.

**CABBAGE:** Downy Mildew in Plant Beds—1½ lbs. 2 to 3 times a week beginning when plants emerge.

**CELERY:** Early Blight (Cercospora), Late Blight (Septoria) and Damping-Off— $\frac{3}{4}$  to 1½ lbs. once a week with first application in the plant bed and continuing in the field. Under severe conditions spray every 3 to 5 days. Remove residues by stripping, trimming and washing.

**CHERRIES:** Leaf Spot—Apply 1½ lbs. 3 times at 14-day intervals starting at petal-fall. Spray foliage again immediately after harvest. Brown Rot—1½ to 1½ lbs. at pink bud stage, full-bloom and petal-fall and at 7 to 10-day intervals starting 2 to 3 weeks before harvest. A spreader may be added to late sprays to avoid blotching fruit.

**CITRUS (Florida):** Scab—Apply 1¼ lbs. at prebloom and  $\frac{2}{3}$  petal-fall and late summer or early fall if heavy flush of growth occurs.

**CRANBERRIES:** Fruit Rots—2¼ lbs. 3 times at 14-day intervals with first spray at mid-bloom. Fairy Ring Disease—7½ lbs. in the fall immediately after harvest. Apply 1 gallon per sq. ft. to the diseased area and up to 3 feet outside of the advancing ring of dead plants. Do not apply later than 28 days after mid-bloom.

**GRAPES:** Black Rot—1½ lbs. just before bloom, immediately after bloom and 14 days after bloom. Additional applications may be made on recommendation of State Experiment Station or State Extension Service authorities at delayed dormant, cluster break, berry touch and 10 to 14 days after berry touch stages. Do not apply later than 7 days before harvest.

**LETTUCE:** Damping-Off in Plant Beds— $\frac{3}{4}$  lb. as a drench (4 to 5 gals. per 100 sq. ft.) when plants emerge and at 14-day intervals.

**PEACHES:** Peach Leaf Curl, Shot-Hole—1½ to 2¼ lbs. at dormant and popcorn. Brown Rot Blossom—1½ to 1½ lbs. at pink bud and 25 percent to 75 percent of full bloom. Fusicoccum Canker—1½ lbs. at 7 to 10-day intervals before bloom and after harvest. Do not apply within 21 days of harvest.

**PEARS:** Scab—1½ lbs. in pink, calyx, first and second-cover sprays. Fabraea, Sooty Blotch— $\frac{3}{4}$  lb. in summer sprays. Do not apply later than 7 days before harvest.

**PINE SEEDLINGS:** Fusiform Rust in Plant Beds—1½ lbs. with suitable spreading agent. Apply at 3 to 7-day intervals from plant emergence as needed.

**PLUMS, PRUNES:** Brown Rot—1½ to 1½ lbs. at popcorn, full-bloom and 2 weeks later. Apply again 5 weeks before harvest and repeat at 7-day intervals during wet weather. Leaf Spot— $\frac{3}{4}$  to 1½ lbs. in fall after harvest and before leaf drop. Repeat in spring as buds swell. Do not apply later than 7 days before harvest.

**RASPBERRIES, BLACKBERRIES, BOYSENBERRIES:** Anthracnose, Rust, Spur Blight—2¼ lbs. delayed dormant, 1½ to 2 lbs. when new canes are 10 to 12 inches high and before bloom. Do not apply later than 40 days before harvest.

**TOBACCO:** Blue Mold— $\frac{2}{3}$  to 3 lbs. at 3 to 4-day intervals beginning when leaves are size of a dime and continue until transplanting time. Use 3 gallons of suspension per 100 sq. yds. of bed when plants are small, increasing to 6 gallons as plants enlarge.

**TOMATOES:** Gray Mold, Sclerotinia (Florida)—1½ to 3 lbs. at 3 to 5-day intervals beginning at transplanting time. Apply in combination with other fungicides used for control of other diseases.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the Chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY guarantee or representation, express or implied concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

Distributed by  
**CHEVRON CHEMICAL COMPANY**

FORM 7524-B

USDA Reg. No. 1965-51

ORTHO DIVISION — San Francisco, California 94120

Product No. 1000

Made in U. S. A.

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRESNO, CALIFORNIA

DES MOINES, IOWA

CHERRY HILL, NEW JERSEY

DALLAS, TEXAS

WHITTIER, CALIFORNIA

HONOLULU, HAWAII



ORLANDO, FLORIDA