

Right Panel

**GENERAL INFORMATION:** PreveX is a completely water miscible fungicide concentrate. PreveX is intended for use on turfgrass which is subject to Pythium Blight (Cottony Blight, Grease Spot). It may be used after germination to control Pythium damping-off on turf, or on established turf to prevent or control Pythium Blight. PreveX does not afford broad spectrum disease control for *Rhizoctonia* spp. (Brown Patch), *Sclerotinia* spp. (Dollar Spot), or *Helminthosporium* spp. (Leaf Spot). Use a broad spectrum fungicide if these diseases coexist with Pythium Blight.

PreveX is also recommended for prevention of root rot and damping-off caused by *Pythium* spp. and *Phytophthora* spp. at all stages of ornamental propagation.

**COMPATIBILITY:** PreveX is compatible with Acti-dione RZ<sup>®</sup>, Acti-dione <sup>®</sup>, Thiram<sup>®</sup>, Acti-dione TGF<sup>®</sup>, and these Acti-dione products combined with ferrous sulfate when used at the recommended rate. PreveX is also compatible with iprodione, ethazol, chloroneb, Proxol<sup>®</sup> 80SP<sup>®</sup> (trichlorfon), thiadimefon, chlorothalonil, diazinon, chlorpyrifos, and thiram.

<sup>®</sup>Trademark of The Upjohn Company

For control of other soil borne diseases such as *Rhizoctonia* spp. or *Fusarium* spp., PreveX can be tank mixed with benomyl or thiofanate-methyl. Use the full rate of each fungicide and follow all directions and restrictions appearing on the labels of these products.

**DIRECTIONS FOR USE:** It is a violation of Federal Law to use this product in a manner inconsistent with its labeling:

<u>TURFGRASS</u>	<u>AREA</u>	<u>Fluid Ounces PREVEX</u>	<u>REMARKS</u>
For Control of Pythium Blight and Pythium Damping-Off on Established Turf and Overseeded Areas	1000 sq. ft.	1 1/2 to 4	<u>Established Turf:</u> Appl. as a preventive treatment during periods of high temperature and humidity favorable to the development of Pythium Blight.
		fluid ounces in 2 to 5 gallons of water	
			<u>Overseeded Areas:</u> Apply after germination to control Pythium Damping-Off.
			<u>Retreatment:</u> Retreat at 7-10 day intervals if conditions remain favorable for disease development. Use the lower rate for the shorter interval and the higher rate for the longer interval within the application rate range. Under severe disease pressure conditions, use the highest recommended rate and shortest interval.

**USE PRECAUTIONS:** Do Not graze or feed droppings from treated areas to livestock. Do not enter treated areas until spray deposits has dried.

<u>ORNAMENTALS</u>	<u>Fluid Ounces PREVEX</u>	<u>REMARKS</u>
For Control of Pythium and Phyto- phthora on Orna- mental Plants; African Violet, Ageratum, Alyssum, Aster, Azalea, Browallia, Carna- nation, Celosia, Chrysanthemum, Coleus, Dahlia, Dusty Miller, Easter Lily, Ferns, Geranium, Gloxinia, Impatiens, Lobelia, Marigold, Pansy, Petunia, Poinsettia, Portulaca, Salyia, Snapdragon, Verbena Violet, and Vinca.	3 fl. oz. per 10 gallons of water	Prevex requires no agitation after initial mixing. Stock solutions of Prevex should be used within one week of mixing. Prevex does not cause phytotoxic symptoms on the listed plants when used as directed. Trials on other ornamental species should be conducted before extensive use.
	3 fl. oz. per 10 gallons of water	<u>Seeding:</u> Apply to the seeding area using 3 quarts of solution/10 square feet. Treat before sowing of seeds requiring light and after covering of seeds requiring dark for germination.
	2 fl. oz. per 10 gallons of water	<u>Transplanting (Seedlings):</u> Apply to the transplanting area using 3 quarts of solution/10 square feet before transplanting.
	2 fl. oz. per 10 gallons of water	<u>Transplanting (Cuttings):</u> Drench the cutting bed using 2 gallons of solution/ 10 square feet fore setting.
	2.5 fl. oz. per 10 gallons of water	<u>Potting:</u> Dip bare root plants in the solution before potting. After potting drench each plot at the rate of 3.5 fl. oz. of solution/4 inch pot. Repeat the drench treatment at 3 to 5 week intervals.
woody Ornamentals	2.5 fl. oz. per 10 gallons of water	For control of <i>Phytophthora</i> spp. on woody ornamentals such as azalea, drench each container at the rate of 5 quarts of solution/10 square feet of surface. Repeat the drench treatment at 3 to 4 month intervals.

#### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in original container and keep closed. Store in a cool, dry place.  
**PESTICIDE DISPOSAL:** wastes resulting from the use of this product may be disposed  
of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or  
reconditioning, or puncture and dispose of in a sanitary landfill, or incineration,  
or, if allowed by State and local authorities, by burning. If burned, stay out of  
smoke.

**IMPORTANT- READ BEFORE USE -** By using this product user accepts the following  
conditions, warranty, disclaimer of warranties, and limitations of liability:  
**CONDITIONS:** The directions for use of this product are believed to be adequate and  
should be followed carefully. However, because of extreme weather and soil  
conditions, manner of use and other factors beyond NOR-AM's control, it is  
impossible for NOR-AM to eliminate all risks associated with use of this product.  
As a result, crop injury or ineffectiveness is always possible. **WARRANTY AND DIS-  
CLAIMER OF WARRANTIES:** NOR-AM warrants that this material conforms to the chemical  
description on the label and is reasonably fit for use under the directions and  
conditions on this label. NOR-AM AND SELLER MAKE NO OTHER EXPRESS OR IMPLIED  
**WARRANTY OF FITNESS, MERCHANTABILITY OR OTHERWISE. LIMITATIONS OF LIABILITY:** The  
liability of NOR-AM or Seller for damages arising from use of this product is  
limited to the replacement cost of the product used and shall not include any conse-  
quential damages such as loss of profits or other values. **NO CHANGES AUTHORIZED:**  
No one other than an authorized Agent to NOR-AM is authorized to make any other  
warranty, or change the above conditions, disclaimer or limitations and then only if  
in writing and with a specific reference to this shell. Product identical to  
PREVEX 3, a registered product of NOR-AM Agricultural Products, Inc. PREVEX 3 is a  
trademark of Geheing AG Berlin-Bergkamen.





Left Panel

PREVEX

TURF AND ORNAMENTAL FUNGICIDE  
For Phythium and Phytonthora Control

Active Ingredient	
Propamocarb hydrochloride.....	66.5%
*Propyl[3-(dimethylamino)propyl]carbamate monohydrochloride	
Inert Ingredients.....	33.5%
	Total 100.0%

Contains 6.0 lbs. active ingredient per gallon.

Keep Out of Reach of Children  
CAUTION-CAUTION

PRECAUCIONAL USUARIO: No usee no see  
ng es. no use este producto hasta que la  
atiqueta le haya sido expuesta ampliamente.

STATEMENT OF PRACTICAL TREATMENT: ...  
wash with soap and water. ...  
eyes. ...  
attention if irritation persists.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: May be harmful if  
swallowed. ...  
ENVIRONMENTAL HAZARDS: Do not apply directly to water or  
wet lands.

SEE REVERSE PANEL FOR  
ADDITIONAL PRECAUTIONS

NET CONTENTS: 1 QUART

EPA Reg. No. 2134-101  
EPA Est. No.

USE BEFORE

U.S. Patent No. 2,543,074

MANUFACTURED FOR  
NOR-AM Agricultural Products, Inc.  
Naperville, Illinois 60560

REV. 10/11/68