

1369-48

### DIRECTIONS

Insert Taps in Top of Can with Sharp Nail

PARSONS INSECTICIDE DUST may be applied at any time when weather conditions are favorable (see other label). The DUST may be applied during the daytime to either moist or dry foliage. Apply DUST regularly at 7 to 10 day intervals, beginning soon after plants are above ground.

Dosage may vary depending on cultural practices, amounts of foliage, soil equipment, etc. For best results keep foliage evenly coated with the DUST. This is especially important on new growth. Do not apply Parsons 1% Rotenone dust to crops listed below less than one day of Harvest.

#### CUCUMBERS, MELONS, SQUASH, PUMPKINS

To control Striped and Twelve Spotted Cucumber Beetles, apply when insects first appear and continue applications weekly until danger of insect injury is past. Redust after rains and continue with regular applications of Parsons Insecticide Dust.

#### TOMATOES

To control Potato Flea Beetles, Colorado Potato Beetles and Small Tomato Hornworms, begin application 2 or 3 days after plants are set in the field. To control Tomato Worms apply when worms are still small. Repeat as necessary.

#### POTATOES and EGG PLANTS

To control Colorado Beetles and Potato Flea Beetles begin application when insects make appearance. Repeat at 5 to 7 day intervals as necessary when danger of insect injury is past. Continue regular applications of Parsons Insecticide Dust.

#### BEANS

Mexican Bean Beetles and Bean Thrips, Bean Leaf Beetles: Start when insect and disease pests first appear. Apply to the underside of the leaves as well as top side. Repeat treatment as often as necessary.

#### CABBAGE

Cabbage Looper Worms, Southern Cabbage Worms, Cross-Striped Cabbage Worms, Imported Cabbageworm. Apply DUST when worms first appear. Repeat as often as necessary.

#### ASPARAGUS

Asparagus Beetle and Spotted Asparagus Beetle: The spears should be dusted with Parsons Insecticide Dust and when possible apply when the insects are predominately in the larval state. Repeat as often as necessary.

# Parsons

### CAUTION

Avoid inhalation of dust. Keep this package away from excessive heat. Avoid storage near steam pipes and radiators.

**ENVIRONMENTAL CAUTIONS**  
THIS PRODUCT WILL KILL FISH.

**KEEP OUT OF ANY BODY OF WATER.**

Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**Do not reuse container. Destroy when empty.**

**NOTICE TO PURCHASER** — Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of Parsons Chemical Works, Inc. and the Seller. Buyer assumes all risks of use; storage or handling of this material not in strict accordance with directions given herewith.

NET WT. NO. 100-41-1

EPA REG. NO. 3369-48

# Parsons

## INSECTICIDE DUST ROTENONE

### 1%

### Controls

## Worms and Insects

See specific names listed on directions panel on

## Vegetables

### CAUTION

**KEEP OUT OF REACH OF CHILDREN**  
See Side Panel for Additional Cautions

#### NET WEIGHTS:

Rotenone .....	1.1%
Other Active Ingredients .....	2.1%
Inert Ingredients .....	97.0%

*Manufactured Since 1916*

PARSONS CHEMICAL WORKS, INC.