



FOR CEREAL LEAF BEETLE

PROGRAM — The Federal Grain Inspection Service

For application to stored grain, apply Malathion 57% Emulsifiable Insecticide to the grain surface. Malathion 57% Emulsifiable Insecticide may be used for control of Cereal Leaf Beetle, Red Flour Beetle, Indian Meal Moth, Confused Flour Beetle, Rice Weevil, Flat Grain Beetle, Rusty Grain Beetle, Lesser Grain Borer, Granary Weevil, and Saw-toothed Grain Beetle. For control of Cereal Leaf Beetle, apply Malathion 57% Emulsifiable Insecticide to the grain surface after the first week of June.

Please note: Malathion 57% Emulsifiable Insecticide should be applied to stored grain in a well-ventilated area. Do not apply to grain which is being stored in a structure which is not well-ventilated. Do not apply to grain which is being stored in a structure which is not well-ventilated.

MATERIALS — Amount of Malathion 57% Emulsifiable Insecticide to be applied to stored grain is given by the following table.

Do not apply to crops within 30 days of harvest.


Sprays Applied by Ground Equipment — Do not apply to stored grain. Do not apply to stored grain. Do not apply to stored grain. Do not apply to stored grain.

WARNINGS — Apply chemical by other than dust from number of days given.

Malathion presents no residual problem from drift onto stored grain or pasture.

Note — When used in aeration of stored grain, should be applied at a rate of 1.0 to 2.0 pounds per acre. Do not apply to stored grain which is being stored in a structure which is not well-ventilated.

1969-96



**MALATHION 57%
PREMIUM GRADE
EMULSIFIABLE INSECTICIDE**

**STORED GRAIN PROTECTANT
AND
EMPTY GRAIN BIN SPRAY**

**GRAIN
ELEVATORS**

**FARMER'S
GRANARIES**

CONTROLS: Cereal leaf beetle, Red flour beetle, Indian meal moth, Confused flour beetle, Rice weevil, Flat grain beetle, Rusty grain beetle, Lesser grain borer, Granary weevil, Saw-toothed grain beetle.

Malathion may be used under the requirements of the Khapra beetle quarantine, where water or diesel oil emulsions are indicated, as prescribed by quarantine instructions.

ACTIVE INGREDIENTS:


Malathion*	57%
Xylene	40%
INERT INGREDIENTS	3%
*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate	

CAUTION — KEEP OUT OF REACH OF CHILDREN


(See Other Panel for Additional Cautions)

PARSONS CHEMICAL WORKS, Inc.
Grand Ledge, Michigan 48837

EST. REG. NO. 1969-MI-1
EPA REG. NO. 1969-96
L107409



Parsons Malathion 57% Emulsifiable Insecticide is a registered trademark of Parsons Chemical Works, Inc.



DIRECTIONS

Follow procedure approved by Federal Grain Sanitation Program. Controls certain insects mentioned on front panel except Cereal Leaf Beetle. Directions for Cereal Leaf Beetle Shown on side panel.

As A Spray: For the control of Cereal Leaf Beetle, apply Malathion 57% Emulsifiable Insecticide to the grain surface. Do not apply to stored grain which is being stored in a structure which is not well-ventilated. Do not apply to stored grain which is being stored in a structure which is not well-ventilated.

