

RECOMMEND

ACCEPTED
10-27-72
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND SOILICIDE ACT
FOR ECONOMIC PESTICIDE REGISTERED
ON 10/27/72

Cameraaman's Note
Poor Copy

1980-9
with
Comments

Residual Fly Spray Concentrate
For Use On Dairy or Beef Cattle, in Barns or Stables

An Effective Concentrate for Control of Face Flies, Horn Flies, Stable Flies, and Lice
and Ticks on Beef Cattle, Dairy Cattle, Swine, and Poultry.

An emulsifiable concentrate, mixes
with water or oil to form easy-to-use
dilutions for direct application. Kills
insects quickly by contact action.

Contains Clopyralid with Hesperidin.

Directions for Use: See label for
Dilution, Application, and Precautions.
For information on the use of this product,
contact your local Agricultural Extension
Office or the manufacturer.
REGISTERED TRADEMARKS: Clopyralid, Hesperidin.
© 1972, Agricultural Research Service, U.S. Department of Agriculture.

ACCEPTED WITH COMMENTS

ACCEPTED

10-27-76

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROBOTICIDE ACT
FOR ECONOMIC POISON REGISTRY
ED UNDER NO. 1980-7

GENERAL DIRECTIONS

PROPER APPLICATION: Mix 6 to 8 quarts Cidrelex with 19 gallons water; mist directly on dairy or beef cattle once weekly. Repeat application as necessary to maintain control.

For application with conventional hydraulic sprayers or low pressure knapsack sprayers, mix required amount with water and apply in accordance with state and federal recommendations.

For use in back rubbers or face treaters to control horn flies and face flies, mix one gallon Cidrelex with 16 gallons oil. Use light base neutral oil or No. 2 kerosene fuel in engine use with local recommendations. Face treaters (sheet canvas type) should be placed in doors, lanes or runways where cattle pass through to water in feeding areas around salt licks where animals may freely treat themselves.

For horn flies, face flies, stable flies, lice and the Lone Star tick on dairy cattle and beef cattle, mix one gallon Cidrelex Insecticide Concentrate with 13 to 16 gallons of water, spray to thoroughly cover all parts of the animal using 2.4 quarts of diluted spray per large animal and proportionately less for smaller animals. Repeat as necessary to maintain control, but do not apply more than once per week. Do not contaminate feed or feed.

As an aid in reducing populations of house flies and stable flies in barns and sheds, on walls, eaves, fences, in pig sheds and corrals, mix 1 gallon Cidrelex with 13 to 20 gallons of water, spray at the rate of one gallon of diluted spray per 1000 square feet of surface to be treated. Repeat as necessary to maintain control.

(Cont'd on right hand panel)

with comments

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

For application in fall and winter on dairy cattle, beef cattle, sheep, goats and hogs, mix one gallon Cidrelex Insecticide Concentrate with 52 to 64 gallons of water and thoroughly apply to cattle at the rate of 3 to 5 gallons of diluted spray per animal. Use proportionately less for smaller animals. Repeat applications as necessary to maintain control.

Toxic to Fish and Wildlife: Keep out of lakes, streams or ponds. Rinse equipment and containers and dispose of liquid wastes in a pit in non-crop land away from water supplies or other surface water. Apply this product only as specified on this label.

Do not contaminate feed or foodstuffs. During commercial or prolonged exposure in spray mix or the loading of containers, wear clean synthetic rubber gloves and a mask or respirator of a type approved by U.S. Bureau of Mines and recommended by U.S. Department of Agriculture for protection against Cidrelex Insecticide. Use material only for recommended purposes as recommended on label.

Caution: This product is highly flammable. Avoid contact with skin, eyes, nose, mouth, and clothing. Wash thoroughly with plenty of water immediately after contact. Do not reuse container when empty.

Auton Equipment Mfg. Co.

Camerman's Note
Poor Copy

ACCEPTED WITH COMMENTS