DANGER

Sanco SANI-POOL Algicide is formulated to compliment swimming pool water being treated with normal chlorine systems. It is especially effective against growth of algae and, being non-volatile, aids in maintaining pool water clarity and sparkle. The residual effectiveness of Sanco SANI-POOL Algicide tends to stabilize the chemical treatment system.

Sanco SANI-POOL Algicide is compatable with all chemicals normally used in swimming pool maintenance, however, in it's concentrated form, this chemical should not come in contact with high concentration of chlorine.

DO NOT ADD SANCO SANI-POOL ALGI-CIDE TOGETHER WITH CHLORINE PRO-

DUCTS.
When adding to the pool water, these chemicals should be added separately. Sanco SANI—POOL. Algacide is economical, efficient & sta-

DANGER

KEEP OUT OF REACH OF CHILDREN

Corrosive. Causes severe eye damage and sirritation. Do not get in eyes, on skin or on clothing. Protect eyes with goggles or face shield when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

ble.

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If Swallowed, drink PROMPTLY a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against cir-



SANI-POOL

ALGICIDE

ACTIVE INGREDIENTS

n- alkyl (50% C₁₄, 40% C₁₂, 10% C₁₆,) dimethyl benzyl ammonium chloride - - - -10.0% INERT INGREDIENTS

DANGER — KEEP OUT OF REACH OF CHILDREN. See antidote statement and additional precautions on side panel. Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

This product is toxic to fish. Treated effluents should not be discharged where it will drain into lakes, streams, ponds or public waters. Apply this product only as specified on this label.

SANCO PRODUCTS COMPANY

DIVISION OF

SANITARY SUPPLY & CHEMICAL CO., Inc.
MACON GA ALBANY GA

culatory shock, respiratory depression and convulsion may be needed.

DIRECTIONS

For an initial application or when pool water is changed, use one gallon Sanco SANI-POOL Algicide per 25,000 gallons of pool water. For maintenance, use four ounces per 5,000 gallons of pool water. The appropriate amount of Sanco SANI-POOL Algicide should be added in small increments directly to pool water while walking around the pool. In this manner, more adequate distribution can be achieved.

If pool has visible algae growth, treat with initial dose. Vacuum pool after 24 hours to remove algae debris. If visible algae are still present or reappear, repeat to atment procedure as necessary. When pool is free of visible algae, use recommended maintenance dose, at five to seven day intervals. During periods of extreme ly high temperatures, heavy use, or make up water added at rate of 10% per week, use maintenance procedure daily; otherwise use maintenance procedure on a five to seven day schedule.

DANGER

Full strength SANI—POOL Algicide causes severe eye and skin irritation. See opposite pannel for further precautionary instructions.

TO DETERMINE POOL CAPACITY:

Rectangular pools: length times width times average depth in feet times 7.5 equals gallors water in pool.

Round and Oval Pools: long diameter times short diameter times average depth times 5.9 equals gallons water in pool.

NET CONTENTS: 1 GALLON

EPA - REG. NO. - 1989 - 1 44 EPA - EST - 1989 - GA - 1

