



**DIRECTIONS**

**LICE, TICKS, SCABIES, SARCOPTIC AND PSOROPTIC MANGE on Hogs, Horses, Sheep and Goats:**

1½ pounds per 100 gallons of water. Apply a second spray two weeks later. Use as a spray or dip, but do not apply to calves under 3 months old. Apply 2 quarts of solution per adult animal; proportionally less on smaller animals. Do not spray within 30 days of slaughter, or dip within 60 days of slaughter. Do not use on dairy goats.

Do not treat animals under 3 months of age. Do not treat sows within 2 weeks of farrowing, nor for at least 3 weeks thereafter. Avoid treating animals in cold, stormy weather. Avoid treating overheated or sick animals.

**LICE ON BEEF CATTLE:**

1 pound per 100 gallons of water, applying 2 quarts of solution per adult animal. Do not apply to animals being finished for slaughter.

Do not apply within 30 days of slaughter.

**SARCOPTIC AND PSOROPTIC MANGE ON BEEF CATTLE:**

1½ pounds per 100 gallons of water. Make a single application of 2 quarts of solution per adult animal; proportionally less on smaller animals. Do not apply to animals being finished for slaughter.

Do not apply within 30 days of slaughter.

**FLY CONTROL: (Residual) FOR BARNs, premises and breeding places:**

Use 10 pounds per 100 gallons of water. Spray thoroughly at the rate of 1 gallon of solution per 500 square feet. Use only as a spot treatment in the home, around doors, screens, and surfaces where flies alight or congregate. Do not use in poultry houses. Do not contaminate feed or drinking water. Do not use in dairy barns.

**HORNFLY CONTROL ON LISTED ANIMALS: (Except Dairy Animals):**

Mix 1 pound per 100 gallons of water, using 2 quarts of solution per adult animal. Repeat as necessary. Do not spray within 30 days of slaughter.


**CONTROL OF INSECTS ON VEGETABLES: Use amount indicated in 100 gallons of water.**

CONTENTS ONE POUND

18-1201



6-10-71  
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 1990-8



**KILLS Lice, Ticks, Sarcoptic and Psoroptic Mange and Scabies, Flies, and Certain Insects on Vegetables.**

<b>ACTIVE INGREDIENTS:</b>	
Gamma Isomer of Benzene Hexachloride (from Lindane) .....	25%
<b>INERT INGREDIENTS</b> .....	75%
<b>TOTAL</b> .....	100%

**WARNING**

Keep out of reach of children  
See side panel for additional precautionary labeling

U.S.D.A. Reg. No. 1990-8

**DIRECTIONS**

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS:**

Aphis, Flea Beetle, Caterpillars, and Thrips use 1 pound per 100 gallons of water or apply at the rate of 3 pounds per acre as an overall treatment. Do not apply after edible parts start to form.

**CUCUMBERS:**

Cucumber Beetle and Thrips use ½ pound per 100 gallons water. Squash Bug, Leaf Miners, Grasshoppers, Lygus Bug, Aphis, Melon Worms, use 1 pound per 100 gallons of water. Use 3 pounds per acre as an overall treatment. Do not apply after blossoms open. Treat under the leaves also. Quality of crop may be reduced due to treatment.

**PEPPERS:**

Aphis, Leaf Miners, Army Worms use 1 pound per 100 gallons of water or 2 4/10 lbs. per acre as an overall treatment. Do not apply after edible parts start to form.

**SPINACH:**

Aphis and Leaf Miners use 1 pound per 100 gallons of water or 3 pounds per acre as an overall treatment. Do not apply after seedling stage.

**STRAWBERRIES:**

Wasp, Spittle Bug use ½ pound per 100 gallons water or ½ pound per acre as an overall treatment. Do not apply after fruit starts to form.

**TOMATOES:**

(For fresh fruit market only.) Aphis, Thrips use ½ pound per 100 gallons of water or 3 pounds per acre as an overall treatment. Do not apply after blossoms fall.

**DIRECTIONS — Seed Treatment**

Dust dry seeds at the following dosages per 100 pounds of seed: Beans, Gladiolas and Daffodils: 2 to 4 ounces; Lettuce, Peas, Carrots, C-ions, Grains and Milo: 4 ounces; Cotton: 8 ounces; Corn: 4 ounces (2¼ ounces per bushel); Cucumbers: 4 ounces; Tomatoes: 8 ounces. If soil temperature is extremely high due to low moisture or very hot weather, reduce dosage by one-half. Seed treatment will afford temporary protection of seed and seedlings from attack by wire-worms, seed corn maggot, kafir ants and onion maggots. Treat seeds shortly before planting. Treated seeds exposed on soil surface will be hazardous to birds and wild life.

**WARNING**

May be fatal if swallowed.  
Harmful vapor and dust.  
Do not breathe the dust or spray mist. Do not get in eyes, or on skin.  
Flush with water, or wash skin with soap and water.  
Avoid contamination of food.  
May produce irritation of skin and eyes.  
May be absorbed through skin. Wash thoroughly after handling. Lindane 25% WP should not be applied to food crops after heads, pods, bunches of blossoms form, as the case may be. Do not apply Lindane 25% WP to crops in rotation with root crops including peanuts.  
Do not spray calves under 3 months old.  
Treated seeds should not be used for feed, food, or oil purposes.  
This product is poisonous to fish and wild life. Keep out of lakes, ponds, and streams. Do not apply in any manner not specified on the label.

NON WARRANTY Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

**FARMLAND INDUSTRIES, INC.**