

Each Quart Is Equivalent to One Pound
50% Wettable Methoxychlor Powder.

18-2404

CHERRIES

For cherry Fruit Fly, use 1 quart in 100 gallons of water. Make first application when flies appear, and repeat applications at 7 to 14-day intervals until harvest. Do not apply within 7 days of harvest.

ALFALFA AND CLOVER

For the control of the alfalfa caterpillar, alfalfa weevil larvae, armyworm, flea beetle, Japanese beetle, leafhoppers, Lygus bug and spittle bug, apply 2 to 3 quarts in 18 to 20 gals. water per acre. Apply when insect populations begin to build up. Do not apply within 7 days of harvest or grazing. For airplane spraying apply 2 to 3 qts. in 3 to 10 gallons water per acre. Use the rate of water that is consistent with crop growth and good coverage.

EMPTY GRAIN BINS

As an aid for control of Rice Weevil, Granary Weevil, Lesser Grain Borer, Angoumois Grain Moth and certain other insect pests of stored grain, during accessible or exposed stages; and as a treatment to help prevent infestation of newly stored grain, apply 1 gallon with 9 gallons of water. Make application to thoroughly cleaned surfaces at the rate of 2 gallons of spray mixture per 1,000 square feet. Allow sprayed surfaces to dry before storing.

COMPATIBILITY

Do not use with materials containing sulfur or other substances incompatible with oil. Do not use on crops within 14 days of application of any material containing sulfur

11-5-68



METHOXYCHLOR
EMULSIFIABLE CONCENTRATE

CONTAINS
2 POUNDS OF METHOXYCHLOR
PER GALLON

ACTIVE INGREDIENTS:

| | |
|--------------------------|------|
| *Methoxychlor, Technical | 24% |
| Methylated Naphthalenes | 63% |
| Dimethyl Phthalate | 10% |
| INERT INGREDIENTS | 3% |
| Total | 100% |

U. S. Patent 2,420,928

*24% Methoxychlor, technical is equivalent to 21%
2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and
3% other isomers and reaction products.

CAUTION

18-2404

11-5-68



METHOXYCHLOR

EMULSIFIABLE CONCENTRATE

CONTAINS
2 POUNDS OF METHOXYCHLOR
PER GALLON

ACTIVE INGREDIENTS:

| | |
|--------------------------------|------|
| *Methoxychlor, Technical | 24% |
| Methylated Naphthalenes | 63% |
| Dimethyl Phthalate | 10% |
| INERT INGREDIENTS | 3% |
| Total | 100% |

U. S. Patent 2,420,028

*24% Methoxychlor, technical is equivalent to 21%
2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and
3% other isomers and reaction products.

CAUTION

U.S.D.A. Reg. No. 199C-16

DIRECTIONS

For use against insects affecting Beef Cattle,
Barns, Warehouses, etc.

For Spraying Beef Cattle for Hornflies, Stable Flies (only supplementary control can be obtained — major control would be in the thorough spraying of barns and treatment of other breeding area) and Lice: Use 1 quart per 12 gallons water, using 2 quarts of the dilute solution per adult animal. Repeat as necessary (2 - 3 weeks).

For Spraying Interior of Barns, etc. For House Flies, Stable Flies and Mosquitoes; use 1/2 gallon per 5 gallons of water. Spray thoroughly walls, ceilings, stalls, screens, manure piles, etc., using 1 gallon dilute spray to each 600-1,000 square feet. Repeat as necessary (4 - 6 weeks).

Do not spray on or near water, ponds, streams, etc. Do not spray on or near crops, fruit, etc. Do not spray on or near birds, etc.

CAUTION Harmful if swallowed. Avoid inhalation of spray mist. May cause skin irritation. Avoid contact with skin. In case of contact, wash with soap and water. Do not use on humans or animals.

Avoid contamination of feed and foodstuffs. Do not store or use near fire or open flame.

Fish will be killed if their waters are contaminated with this product.

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.