

ACCEPTED
 Apr 17, 1972
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PESTICIDE REGISTRY
 REGISTRY NO. 1990-27 COMMENTS

18-2650



CONDITIONS OF SALE

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

CAUTION

Do not use treated seed for flour or feed. While sowing seed, Copper Carbonate dust may sift into the working parts of the drill; therefore, before using it again, free the feed wheels by turning the feed shaft with a wrench to avoid breaking of shaft or gears.

CONTENTS ONE POUND
 U.S.D.A. Reg. No. 1990-27

ACCEPTED WITH COMMENTS



20% COPPER CARBONATE
 Containing 20% Metallic Copper
 FOR TREATING WHEAT AND SORGHUM SEED

ACTIVE INGREDIENTS:

Metallic Copper (in basic copper carbonate)	20%
INERT INGREDIENTS	80%
TOTAL	100%

CAUTION

Keep out of reach of children
 See side panel for additional precautionary labeling

DIRECTIONS FOR USE

For the control of stinking smut of wheat and kernel smut of sorghum, treat clean seed any time up to four weeks before planting. Use three ounces Copper Carbonate 20% per bushel of sorghum, and two to three ounces per bushel of wheat. Seeds to be treated should be placed with the specified amount of Copper Carbonate in a tightly closed container and shaken vigorously for two or three minutes or until each seed is coated lightly with the dust. A rotating drum treater can be used for large quantities of seed. Following treatment the seed should be poured out on a screen to get rid of any excess Copper Carbonate. This excess dust can be used again.

WARNING

Poisonous if taken internally. Irritating if inhaled. Avoid breathing dust. Use a respirator.

COOP