


18-1489



ACCEPTED
Feb-3-1969
1990-80



U.S.D.A. Reg. No. 1990-80

**HEPTACHLOR
3E**

CORN ROOTWORM—Mix Heptachlor 3E at 2 quarts per acre in sufficient water to cover one acre. Apply the spray broadcast to the ground and disc or work into the top 3 or 4 inches of soil. If planter spray attachment is used, it may be applied as a band in the row at 1/3 gal. per acre at time of planting corn.

WIRE WORM IN CORN BROADCAST—Mix 2/3 to 1 gallon Heptachlor 3E with sufficient water to spray one acre. Apply prior to planting, and thoroughly work into top few inches of soil. For row application in corn, mix 1/2 to 3/4 gallon Heptachlor 3E in sufficient water to spray one acre at planting time.

WARNING

May be fatal if swallowed. Do not get in eyes, on skin, or on clothing. Do not breathe vapor, fumes, or spray mist. Do not store with or in conjunction with feed and foodstuffs. This product is toxic to fish and wild life. Keep out of lakes, streams, and ponds. Apply this product only as specified on the label.

ANTIDOTE

EXTERNAL — In case of spillage on skin, wash promptly with soap and water.
INTERNAL — Emetic of tablespoonful of mustard in a glass of water. Call a physician.

IMPORTANT

Do not reuse empty container. Destroy and bury in a safe place.

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

FARM AND INDUSTRIES

CONTAINS 3 POUNDS HEPTACHLOR 3E PER GALLON

ACTIVE INGREDIENTS:

Heptachlor*	31.00%
Related Compounds	12.10%
Petroleum Distillate	50.90%
INERT INGREDIENTS	6.00%
TOTAL	100.00%

*Heptachlorotetrahydro-4, 7-methanoindene

WARNING

Keep out of reach of children
See side panel for additional precautionary directions