


Code No. 18-1115

**DIRECTIONS**



**FOR HORNFY CONTROL ON DAIRY CATTLE:**

Use 3 to 5 tablespoonfuls per animal. Dust on poll, neck, back, and upper sides of the animal. Rub in lightly to insure that the material is worked down beneath the hair. Repeat application once every three weeks.

For hornfly, lice and flea control on beef cattle and hogs, apply 5 to 10 tablespoonfuls per animal. Rub in lightly to insure that the material is worked down beneath the hair. Provide thorough coverage of the animal.

CONTAINS ONE POUND

**DAIRY DUST**

**A POWDER CONTROL FOR HORNFLIES, LICE, AND FLEAS**

**ACTIVE INGREDIENTS:**

*Technical Methoxychlor .....	10.99%
**Technical Piperonyl Butoxide .....	0.60%
Pyrethrins .....	0.06%
INERT INGREDIENTS .....	88.35%
Total .....	100%

\*Equivalent to 9.67% of 2,2-bis (p-methoxyphenyl) -1, 1, 1-trichloroethane and 1.32% of other isomers and related compounds.

\*\*Equivalent to 0.48% (butylcarbityl) (6-propylpiperonyl) ether and 0.12% related compounds.

**CAUTION**

**Keep Out of Reach of Children**

Code No. 18-1115



# DAIRY DUST

A POWDER CONTROL FOR HORNFLIES, LICE,  
AND FLEAS

ACTIVE INGREDIENTS:

*Technical Methoxychlor .....	10.99%
**Technical Piperonyl Butoxide .....	0.60%
Pyrethrins .....	0.06%
INERT INGREDIENTS .....	88.35%
Total .....	100%

\*Equivalent to 9.67% of 2,2-bis (p-methoxyphenyl) -1, 1, 1-trichloroethane and 1.32% of other isomers and related compounds.

\*\*Equivalent to 0.48% (butylcarbityl) (6-propylpiperonyl) ether and 0.12% related compounds.

**CAUTION**


Keep Out of Reach of Children

U.S.D.A. Reg. No. 1990-91

**FOR SHEEP:**

Direct dusting after shearing will give temporary relief from ticks (keds). Use same method for treating as recommended for hornfly control on dairy animals.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



FARMLAND INDUSTRIES, INC.