



Code No. 18-1222


Chlordane is powerful and effective in controlling insects — by contact and when eaten.

#### USES

To control ants, chinch bugs and chiggers, dust lightly at the rate of 1 pound to 500 to 600 sq. ft. of area. For white grubs, Japanese beetle grubs and sod webworm (lawn moth) apply liberally at the rate of 1 pound to 250 to 300 sq. ft. Water well with hose to get material into the soil. Effective entire season.

To control wireworms, white grubs, Japanese beetle grubs, cabbage maggots and seed corn maggots on radishes, onions and cabbage, work into top 2 or 3 inches of soil when preparing seed beds. One pound treats 600 to 700 square feet. Remains effective all season.

To control cutworms, earwigs, mole crickets, sowbugs, millipedes, and leaf cutting ants on tomatoes, cabbage, peas, and beans, dust liberally at the time of planting or transplanting on the ground around the plants. Remains effective at least 10 days. One pound treats 250 to 300 square feet.



## 6% CHLORDANE DUST

for ants, cutworms, chinch bugs, white grubs, wireworms, lawn moths, cabbage maggots and seed corn maggots infesting lawn, garden and soil

#### ACTIVE INGREDIENTS:

Technical Chlordane\* ..... 6%  
INERT INGREDIENTS ..... 94%  
TOTAL ..... 100%

\*Equivalent to 3.6% octachloro-4,7-Methanotetrahydroindane and 2.4% related compounds.



Code No. 18-1222



## DIRECTIONS

Powerful and effective in  
— by contact and when

## USES

For ants, chinch bugs and  
at the rate of 1 pound  
of area. For white grubs,  
grubs and sod webworm  
liberally at the rate of  
300 sq. ft. Water well  
material into the soil. Effic-

to control wireworms,  
these beetle grubs, cab-  
seed corn maggots on  
and cabbage, work into  
of soil when preparing  
pound treats 600 to 700  
ns effective all season.  
rol cutworms, earwigs,  
vbugs, millipedes, and  
on tomatoes, cabbage,  
ust liberally at the time  
planting on the ground  
. Remains effective at  
pound treats 250 to 300

**6% CHLORDANE  
DUST**

for ants, cutworms, chinch bugs, white grubs, wire-  
worms, lawn moths, cabbage maggots and seed corn  
maggots infesting lawn, garden and soil.

## ACTIVE INGREDIENTS:

Technical Chlordane	6%	*Equivalent to 3.6% octachloro-4,7-
INERT INGREDIENTS	94%	Methanotetrahydroindane and 2.4%
TOTAL	100%	related compounds.

## CAUTION

KEEP OUT OF THE REACH OF CHILDREN

E.P.A. Reg. No. 1990-117

Foliage Dust: To control grasshoppers,  
thrips, and plant bugs on flowers, shrubs  
and other ornamentals, dust directly on the  
plant and insect. Repeat as necessary to  
prevent damage.

## CAUTION

This material, if taken internally, can  
cause toxic symptoms. Do not breathe dust.  
Do not contaminate feed or foodstuffs.

Birds feeding on treated areas may be  
harmful. Fish may be killed if their waters  
are contaminated with this product.

Buyer assumes all risks of use, storage or  
handling of this material not in accord-  
ance with directions given herewith.

This product is toxic to fish and wildlife.  
Keep out of lakes, streams, or ponds. Do  
not apply when weather conditions favor  
drift from areas treated. Do not contami-  
nate water by cleaning of equipment or  
disposal of wastes.

Do not reuse container. Destroy when  
empty.

Keep children and pets off treated areas  
until this material is worked into the soil  
and grass is dry.