LAWN

Maintain a healthy and vigorous growing lawn. Fertilize twice a year. The first application should be made in the spring about April 1 at the rate of 2 pounds per 100 square feet. The second application should be made in the fall about September 1 at the rate of 2 pounds of material per 100 square feet. Apply the material with a fertilizer spreader or by hand as a broadcast treatment. Do not apply when the grass is wet as possible grass burn may occur. Keep children and pets off treated areas until material has been thoroughly watered into the soil and the grass has completely dried. The following lawn pests are controlled at the above rates of application: ants, army worms, chiggers, Japanese beetle larvae, ~~sod webworms~~, white fringed beetle larvae, white grubs, and wire worms.

RENOVATING OLD LAWNS - ESTABLISHING NEW LAWNS

Apply Coop Big 2 at the rate of 3 to 5 pounds per 100 square feet and work well into the upper 2 to 4 inches of soil, prior to seeding. When the new seedlings are rapidly growing in the spring, a second follow-up application at the rate of 2 pounds per 100 square feet, should be applied.

SHRUBS

Apply Coop Big 2 in the spring when the shrubs are in bud. Loosen the soil around the shrub with a hoe and mix in a heaping handful of fertilizer. Water the plant well after treatment. Shrubs should not be fertilized in the fall as growth will be encouraged and may result in winter damage to the plant.

FRUIT AND SHADE TREES

Apply Coop Big 2 around the drip line of the tree at the rate of 1/4 to 1/2 pound per inch of the tree's circumference. After 2 or 3 years when the tree is well established, holes approximately 2 inches in diameter and 12 to 24 inches deep may be dug around the drip line of the tree and the fertilizer distributed evenly in the holes at the above rates. Water the tree well after applying the fertilizer.

VEGETABLE GARDEN

Apply Coop Big 2 at the rate of 1 to 2 pounds per 100 square feet prior to spading or plowing and work well into the soil. A single application will control armyworms, Japanese beetle larvae, white-fringed beetle larvae, white grubs, and wireworms attacking cauliflower, cabbage, cantaloupe, lettuce, peppers, radishes, tomatoes, and turnips.

FLOWER GARDEN

Apply Coop Big 2 at the rate of 1 to 2 pounds per 100 square feet to the soil surface prior to the seed bed preparation for annual flowers and prior to setting out perennial plants. Established perennials should be treated as directed for shrubs.

Avoid contaminating food. Avoid contamination of food and feedstuffs.

. . . pounds net

. . . .

Controls insects . . . . lawn and garden.

Contains .7% Chlordane.

ACTIVE INGREDIENTS
Technical Chlordane*   .7

INERT INGREDIENTS   99.3

TOTAL   100.0

*Equivalent to .4% octachloro 4,7 methane tetrahydroindane and .3% related compounds.

FERTILIZER - GUARANTEED ANALYSIS

Nitrogen   11
Available Phosphoric Acid   11
Available Potash   11

CAUTION (18 point type)

KEEP OUT OF REACH OF CHILDREN (18 point type)

Manufactured for and distributed by
Farmland Industries
Kansas City, Missouri

Cyclone Lawn Spreaders:

Rate	Micro-Min. Setting Once Over Twice Over	Spread Width
2 pounds per 100 square feet	8.5	4.5	6 feet

Imperial Lawn Spreaders:

Model No.	Rate	Spreader Setting
. . . . . .	2 lbs. per 100 sq. ft.	9
. . . . Imperial	same as above	8-3/4
. . . . Professional	same as above	8
. . . . Husky	same as above	8-3/4

O. M. Scott Lawn Spreaders:

Model No.	Rate	Spreader Setting
. . . .	2 lbs. per 100 sq. ft.	8

The Cyclone Lawn spreader's feature - taper edge spread - will not . . . . the "edge of spreading line" on a once-over application. . . . . . for maximum spreading accuracy, the twice-over application is recommended.

. . . twice-over application with other spreaders, use one-half the normal setting.