


U.S.D.A. Reg. No. 1990-153




DRY INSECTICIDE

DIRECTIONS FOR USE

ACCEPTED
Dec-8-1967
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 1990-153 SUBJECT
TO ATTACHED COMMENTS.

Code No. 18-1022



SWINE

For the control of LICE, make thorough application to the animals. In addition, pens and sleeping quarters should be thoroughly cleaned and then generously dusted with DRY INSECTICIDE. Repeat application in 10 days, then thereafter as needed.

POULTRY

For the control of CHICKEN RED MITE and NORTHERN FOWL MITE in poultry houses, apply dust thoroughly to roosts, nests, and adjacent areas and to the floor litter at the rate of 1 pound per 50 - 60 square feet of floor surface. Repeat application as necessary. Do not apply directly to the birds.

BEEF CATTLE

For the control of LICE make a thorough application to the animals. Remove all dirt and filth from inside sleeping quarters and pens. Sprinkle DRY INSECTICIDE generously in housing and pens once or twice each week, or until lice disappear. Repeat application to animals in 10 days and thereafter as needed. DO NOT APPLY TO LACTATING ANIMALS WITHIN 2 WEEKS OF FRESHENING. DO NOT TREAT CALVES UNDER ONE MONTH OF AGE.

DAIRY CATTLE

For the control of HORN FLIES, apply 2 ounces (4 tablespoons) of dust per animal. Apply the dust on the back and neck of the animal and repeat application at 10 - 14 day intervals. Do not apply to lactating dairy cattle later than 5 hours before milking or during milking.

SHEEP and GOATS

For the control of LICE, TICKS and KEDS on sheep and goats, treat animals thoroughly. Repeat application after 2 or 3 weeks if needed. Remove all dirt and filth from inside sleeping quarters and pens. Sprinkle DRY INSECTICIDE generously in housing quarters and pens once or twice each week, or until lice disappear. Do Not Apply to Milk Goats. Do Not Treat Animals Under One Month of Age.

CAUTION

Avoid prolonged breathing of dust or spray mist. Avoid prolonged or repeated contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on the label.

NOTICE

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.

Manufactured for and Distributed by

FARMLAND INDUSTRIES

KANSAS CITY, MISSOURI



ACTIVE INGREDIENTS:

Malathion*	4.00%
Sulphur	17.50%
INERT INGREDIENTS:	78.50%
Total	100.00%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION

Keep out of reach of children
See side panel for additional precautionary labeling

NET WEIGHT 25 POUNDS