
DIRECTIONS

- quarts CO-OP Beef Cattle Spray per 100 gallons of water. Spray or dip animals thoroughly. Do not treat within 30 days of slaughter. Always stir vat before using and following each replenishment. Water animals before dipping to prevent them from drinking dip solution.
- BLEF ANIMALS Horn flies, mosquitoes, gnats, and as an aid in the control of face flies Mix 6 pints CO-OP Beef Cattle Spray per 100 gallons of water. Spray animals thoroughly. Do not treat within 30 days of slaughter.
- Sarcoptic mange and lice. Use 6 pints of CO-OP Beef Cattle Spray in 100 gallons of water. Do not treat within 28 days of slaughter.
- SHEEP Ticks (Keds), Scab (Scabies). Mix 2 quarts CO-OP Beef Cattle Spray per 100 gallons of water. Use as a dip. Animals should remain in dipping vat for 2 minutes. Do not treat within 30 days of slaughter.
- IN CATTLE OILERS for the control of horn flies, mosquitoes and gnats and as an aid in reducing face flies and lice on beef cattle. Use 1 pint of CO-OP Beef Cattle Spray to 2 gallons of oil and saturate the back rubbers at the rate of 1 gallon per 20 feet of cable. Do not apply to animals within 30 days of slaughter. Rubbing units should be placed in areas where animals loaf, feed or water and at a height for cattle to rub their faces (face flies).
- Treatment for scab and mange mites should be made twice with an interval of 10-14 days. Repeat as necessary for lice, ticks, and horn flies.

E.P.A. Est. No. 32761-MO-2

CONTENTS ONE GALLON

18-1890





[TR 96

Under the Federal
For the Federal
and the region of the first second of the first seco

BEEF CATTLE SPRAY

EMULSIFIABLE CONCENTRATE


Aids in Controlling Horn Flies, Face Flies, Mosquitoes and Gnats Contains 6 Pounds Toxaphene per Gallon 12.5% Emulsifier ACTIVE INGREDIENTS

*Technical chlorinated camphene with 67% to 69% Chlorine.
MIXES WITH WATER IN ALL PROPORTIONS

WARNING

Mouse and fractional production appears to find

18-1890





I TR 9 6 1975

Under the Pederal Insecticide,

For all a sticide Act,

region of the Pederal Insecticide,

region of the Pederal Insecticide,

region of the Pederal Insecticide,

BEEF CATILE SPRAY

EMULSIFIABLE CONCENTRATE

Aids in Controlling Horn Flies, Face Flies, Mosquitoes and Gnats Contains 6 Pounds Toxaphene per Gallon 12.5% Emulsifier ACTIVE INGREDIENTS

*Toxaphene	60.0%
Aliphatic Petroleum Solvent	27.5%
INERT INGREDIENTS	12.5%
Total	400 00/

*Technical chlorinated camphene with 67% to 69% Chlorine.
MIXES WITH WATER IN ALL PROPORTIONS

WARNING

Keep out of reach of children.
See side panel for additional precautionary labeling

E.P.A. Reg. No. 1990-156

WARNING

Do not get in eyes, on skin or clothing: Do not breathe mist from sprays. Use of respirator and goggles is advisable. Keep clothing free from residues. Wash thoroughly after using. Keep away from children, domestic animals and food. May be fatal if swallowed. Do not store near feed and food products.

Do not apply to young calves under 3 months of age. Do not apply to dairy animals.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Avoid treating animals during cold stormy weather and over heated or sick animals.

IMPORTANT

Rinse spray equipment and containers and dispose of liquid wastes in a pit in noncrop lands located away from water supplies. Do not reuse container. Depending on the nature of the container, it should be broken, perforated, or crushed before burying. Do not burn.

DO NOT USE, POUR, SPILL, OF STORE NEAR HEAT OR OPEN FLAME MAY BE FATAL IF SWALLOWED

NONWARRANTY

By yer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Francisco Carlo Ca