

ACCEPTED
APR 14 1975
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, or the pesticide registration under EPA Reg. No.



DIRECTIONS FOR USE

Dairy Barn and Dairy Animal Use

HORN FLY CONTROL ON DAIRY ANIMALS: Apply 1 heaping tablespoonful of 50% Methoxychlor as a dust to the poll, neck, back and upper side of the animal and rub in lightly by hand to cover under beneath the hair. May be repeated every 3 days if necessary.

HOUSE FLIES AND STABLE FLIES IN BARN BUILDINGS — Such as poultry houses, livestock shelters, dairy and hog barns. Use 1 lb. in 2 1/2 gallons water. Apply to all surfaces on which flies can crawl. Repeat applications as infestation warrants. Remove animals and protect feed and water from contamination by spray. Use 1 gallon of diluted insecticide per 500 square feet of surface area. Supplement treatment by using proper sanitary practices.



CATTLE LICE ON BEEF CATTLE, HOG LICE ON HOGS AND SHEEP TICKS ON SHEEP: Use 1 pound per 10-12 gallons of water. Spray to get thorough coverage of the animal, using 2 quarts per hog or sheep and 4 quarts per animal of cattle. Repeat spraying operations in 14-16 day intervals.

VEGETABLE INSECTS: For the control of the insect pests on the plants listed below, use 1/2 pound per 100 gallons of water and apply as a thorough cover spray to upper and lower surfaces of foliage and fruit at a rate of 100-150 gallons per acre. Start treatment at first signs of pest infestation and repeat at 7-10 day intervals as infestation warrants.

- | | |
|-------------------------------|---------------------------------|
| Asparagus Beetle on Asparagus | Japanese Beetle on Beans |
| Blister Beetles on Beans | Leafhopper on Beans and Carrots |
| Cabbageworms on Cabbage | Mexican Bean Beetle on Beans |
| Corn Earworm on Beans | Pea Weevil on Peas |
| Cucumber Beetle on Cucumbers | Squash Vine Borer on Squash |
| Flea Beetles on Beans | Tomato Pinworm on Tomatoes |

E.P.A. Est. No. 32751-M-1

18 1882



ACTIVE INGREDIENTS:
 Technical Methoxychlor* 50%
INERT INGREDIENTS 50%
Total 100%

U.S. Patent No. 2,420,928
 *Equivalent to 44% of 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 6% of other isomers and related compounds.

CAUTION: Remove Spray deposits from plant material consumed as food.

CAUTION
 Keep out of reach of children
 See side panel for additional precautionary labeling

E.P.A. Reg. No. 1900-164

DIRECTIONS FOR USE

Asparagus, cabbage, beans, cucumbers, melons, peas, tomatoes, peppers and squash. Do not apply within 7 days of harvest. Do not apply to beets and root crops within 14 days of harvest.

FRUIT INSECTS — CHERRY FRUIT FLY: Apply 2 to 4 lbs per 100 gallons of water as thorough cover spray, as soon as flies appear. Repeat at 7-10 day intervals as long as flies are present. Consult state extension service specialists for timing of sprays. Do not apply within 7 days of harvest. May be used on peaches, plums, prunes, cherries, apricots and nectarines. Do not treat within 7 days of harvest on plums, prunes and cherries, within 21 days on peaches, apricots and nectarines.

PLUM CURCULIO ON PEACHES, PLUMS AND APPLES: Apply 2 to 4 pounds per 100 gallons of water as thorough cover spray. Make first application at petal fall and 2 additional applications at 7-day intervals. Do not apply within 7 days of harvest on apples and cherries. Do not apply within 21 days of harvest on peaches.

GRAPEBERRY MOTH: On grapes use 2 pounds per acre in 100 gallons of water and make first application as a post-blossom treatment, followed by a second application 10 days later. Apply as thorough cover spray. Make third and fourth applications (if necessary) according to treatment schedule recommended by state experiment station. Do not apply within 14 days of harvest.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

CAUTION: This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not reuse empty container. Destroy it by burning with waste or burning. Stay away from open fires.

FARMLAND INDUSTRIES, INC.

NET WEIGHT ONE POUND