


U.S.D.A. Reg. No. 1990-325

U. S. Patent 2,088,651

Code 18-3302

CONTENTS: FOUR POUNDS ACCEPTED



Aug-14-1969

1990-325

Powdered Standard LEAD ARSENATE

ACTIVE INGREDIENTS:

Dry Lead Arsenate	98%
INERT INGREDIENTS	2%
TOTAL	100%

Total Arsenic expressed as Metallic Arsenic (As) 20.50%

Total Lead, expressed as Metallic Lead (Pb) 58.50%

Arsenic in water soluble form expressed as Metallic Arsenic (As)	25%
---	-----

ANTIDOTE: Internal: Give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Have patient lie down and keep quiet. Call a physician immediately. External: If the skin is contaminated, wash immediately with soap and warm water.



DANGER

POISON



KEEP OUT OF REACH OF CHILDREN.

See antidote statement and other warnings on the back panel

Distributed by

FARMLAND INDUSTRIES

KANSAS CITY, MISSOURI 64116

DIRECTIONS

MIXING: Partly fill the spray tank with water. Start the agitator and gradually add the required amount of Lead Arsenate as the tank is filled.

APPLICATION: Thoroughly wet all exposed plant parts with spray mixture.

APPLES AND PEARS: For codling moth, plum curculio, tent caterpillar and certain other chewing insects, use 1 pound together with 1 pound hydrated lime (spray grade) in 33 gallons water. For small sprayers, use 2 level tablespoons with an equal amount of hydrated lime in 1 gallon water. Apply sprays at petal fall and repeat every 7 to 14 days as long as necessary. Do not apply with 40 days of harvest. Note: Residues remaining at harvest should be removed by washing, wiping, or other effective means.

FOR DUSTING: Thoroughly mix 1 part Lead Arsenate and 7 parts hydrated lime. Apply uniformly as often as necessary to keep plants protected. Dusts may be used on potatoes, tomatoes, cabbage, tobacco, and ornamental plants. Precautions given under the above crop headings should be observed. NOTE: RESIDUES remaining at harvest should be removed by washing, wiping, or other effective means.

This application will control some insects listed under ~~potatoes and~~ tomatoes, ~~cabbage~~, and Tobacco mentioned previously. Follow same cautions.

Do not graze livestock in treated groves or orchards.

POTATOES AND TOMATOES: For horn worm, Colorado potato beetle and certain other chewing insects, use 1 pound in 33 gallons water, or 2 level tablespoons in 1 gallon water. Apply when the insects first appear and repeat as necessary to keep plants protected. Do not apply to tomatoes within 40 days of harvest. NOTE: Residues remaining at harvest should be removed by washing, wiping, or other effective means.

CABBAGE: For cabbage worms use 1/2 level pound in 12 gallons water or 1/2 level tablespoon in 1 gallon water. Apply when the insects first appear and repeat as necessary to keep plants protected. Do not apply after weeds begin to form.

TOBACCO: For horn worm, use 1 pound in 12 gallons water and apply to 1/4 acre. For large plants more than 12 gallons of spray mixture may be needed. Apply spray when insects first appear and repeat as necessary to keep plants protected. Use of insecticides may leave undesirable residues and every precaution should be taken in spraying tobacco. Avoid excessive residues.

SHADE TREES AND ORNAMENTAL PLANTS:

For Japanese beetle and certain other chewing insects use 1/2 to 1 pound in 16 gallons water, or 2 to 4 level tablespoonful in 1 gallon water, with an equal amount of hydrated lime (spray grade). Apply spray when insects first appear and repeat as necessary to keep plants protected. SOIL TREATMENT FOR LAWN AND ORNAMENTAL AREAS: For Japanese beetle grubs, white grubs and certain other grubs, apply 1 pound per 100 square feet of lawn area, either as a spray in water, or dry. Spread or spray uniformly over the area to be treated, then wash insecticide (either dry or spray) off grass blades into soil by spraying lawn with clear water. When preparing lawn for seeding the material may be raked into the soil to a depth of 1 or 2 inches about 30 days before planting grass seed.

CAUTION

Keep children and pets off treated areas.

WARNING: Poisonous if swallowed, inhaled, or absorbed through the skin. Do not get in eyes or on skin. Wear protective clothing, natural synthetic rubber gloves, and goggles. If spilled on skin, remove contaminated clothing and wash skin immediately with soap and warm water. Do not breathe fumes, dust, spray mist. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for protection against lead arsenate. Information as to suitable types of masks or respirators is available from the U. S. Department of Agriculture. Do not contaminate feed or foodstuffs. Keep all unprotected persons out of the operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until the insecticide has dissipated. Keep out of reach of children. Wash hands, arms and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is toxic to wild-life. Do not apply to any area not specified on label. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes or by burning. (Keep out of smoke.)

SYMPTOMS OF POISONING: A sense of "tightness" in the chest. Stomach pains. Vomiting and diarrhea.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith.