

Net weight shown on bag

CO-OP

FLY AND WORM CONTROL MINERAL

Medicated

Flies, horn and face flies in the manure of cattle; aid in the prevention of stomach worms, hookworms and nodular worms in cattle; aid in the prevention of foot rot and other diseases of the hoof in cattle; nutritional source of iodine.

ACTIVE DRUG INGREDIENTS:

Phenothiazine*	.....	0.01%
INACTIVE INGREDIENTS	.....	98.0%
Total	.....	100.0%

ACTIVE DRUG INGREDIENT:

Ethylenediamine Dihydriodide ..... 0.01%

\* Biologically assayed

\*\*Refers only to ingredients which are neither insecticides nor drugs.

GUARANTEED ANALYSIS

Calcium (Ca), not more than	.....	7.0%
Calcium (Ca), not less than	.....	6.0%
Phosphorus (P), not less than	.....	6.0%
Salt (NaCl), not more than	.....	24.0%
Salt (NaCl), not less than	.....	20.0%
Iodine (I), not less than	.....	0.025%

INGREDIENTS

Dicalcium Phosphate, Monocalcium Phosphate, Ground Limestone, Salt, Cane Molasses, Manganese Oxide, Iron Carbonate, Iron Oxide, Zinc Oxide, Copper Oxide, Cobalt Carbonate, Potassium Iodide, Vitamin A Palmitate, D-Activated Animal Sterol (Source of Vitamin D<sub>3</sub>), Corn Distillers Dried Solubles, 8% Soybean Hulls, Anise and Mineral Oil.

FEEDING DIRECTIONS

Self feed continuously to cattle at the rate of 1/2 oz. of CO-OP Fly and Worm Control Mineral, Medicated per day for each 100 lbs. of body weight. Each 1/2 oz. supplies 1/4 gram of phenothiazine. Locate this mineral near animals' water, or near their loafing area. Do not feed any other salt or mineral. For maximum protection against horn flies, cattle should be sprayed at the start of the fly season, and self fed CO-OP Fly and Worm Control Mineral, Medicated continuously, beginning several weeks before the fly season until all adult horn flies have disappeared in the fall. This mineral prevents the breeding of horn flies and face flies in the manure of treated cattle, but does not affect existing flies.

Self feed continuously to sheep and goats so each head will receive 2 oz. daily to provide 1 gram of phenothiazine as an aid in the prevention of stomach worms, hookworms and nodular worms.

WARNING: Do not feed to lactating dairy animals. Occasional sensitivity to phenothiazine has been reported in cattle. KEEP OUT OF THE REACH OF CHILDREN.

ACCEPTED

Manufactured by  
FAIRLAND INDUSTRIES, INC.

General Offices: Kansas City, Mo.

USDA REG. NO. \_\_\_\_\_

