#### KORLAN 24E Livestock Remedy -- Not For Human Use

CHECTIONS IN the L4E moves readily with water or oil. To prepare a finished spray, see chart below.

Type and Use	% Ronnel		int Korlan : o use for: 3 Gal.	24E 25 Gal.	Remarks
LIVESTOCK INSECTS					Retreat as necessary but not oftener
Cattle, Calves, Horn thes Screw worms		212000	8 oz	2 qt	than every 2 weeks Dilute with water only Spray entire animal to wet hair coat thoroughly
Lice	' 4 ° o	1'4 02	4	1 զ։	wer hair toar moroughly
Lone star and winter Ticks	14 00	314 02	12 02	3 q!	
Sheep, Goats, Lambs and Kids Screw worms, wool maggots, hornflies	· . ° •	21: oz	8 ¬z	.* qt	
Lice and sheep keds	1400	1'4 OZ	4 07	1 at	
Hogs lice	1400	114 02	4 0z	1 31	
Cal ther — For hindles Aids tinuously Aids in the control of face flies can rub face. Place where animals travel animals may use backrubbers to within 3 may use backrubbers to within 2 weeks.	if backrub regularly weeks of	bers are pla or congred I freshening	ared so an gate: Dry (	nma) daity	Mix 1 gallon Korlan 24E with 27 gailons diesel or mineral oil Do not mix with used motor oil Saturate liber Recharge as necessary
RESIDUAL FLY SPRAY-FARM A 12% spray will give about 2 weeks control A 1% spray will give 4 - 6 weeks control	ol				Spray walls, ceilings, partitions to obtain complete coverage with no runoff Use 1 gallon of diluted spray per 500 - 1000 sq. ft.
Farm Buildings - Including poultry	1200	2'2% 02	8 oz	2 at	Add 3 ounces of sugar per gallon for
houses, livestock shelters, milk rooms, dairy and hog barns	or 1%	or 5 oz	or 1 pt	or 1 gl	added effectiveness. To prevent con- tamination keep milk and milking equipment covered while spraying
Caged Layers - On wire	2°,	10 oz	, 3,	ان 2	Spray poultry droppings beneath
RESIDUAL FLY SPRAY - GENERAL A 12% spray will give about 2 weeks control A 1% spray will give 4 - 6 weeks control				Spray walls ceilings partitions to obtain complete coverage with no runoff Use 1 gation of diluted spray per 500 - 1000 sq. ft.	
Food Process ng Plants	1,00	2'2 02	8 oz	2 qt	Spray when not in operation Cover
•	or 1°°	5 oz	Or 1 pt	or 1 gl	produce and equipment Spraying outside area will improve control
Dwellings - Outside areas, including	1200	21: 32	8 02	2 q!	Spray to point of runoff
porches, patios garages etc	or 1°°	or 5 oz	1 pt	or 1 gi	
Restaurants, Supermarkets Ethics	1200	21: 07	8 02	2 at	Spray outside walls, receiving docks docrways, and refuse areas
Outside Area	0 <b>f</b> 1°a	01 5 02	or 1 pt	or 1 gi	docrways and refuse areas
Meat Packing Plants	1°.	5 oz	t pt	1 gi	Spray outer and inedible product areas
Pouttry Processing Plants	13.	5 ·z	זק ז	1 gi	Spray outer and inedible product areas
FLY BAIT - Where flies congregates, has floors, windows, doorways, et	?°.	10 oz	4 74	3 g.	Add 3 sunces of sugar per gallon diluted spray. Spray or sprinkle at 1 to 2 day intervals until control is obtained.
COCKROACH CONTROL - Use 1% spray for Germ 2 1					Apply as a spot spray to all areas frequented by cockroaches
Buildings - Including dwellings	••.	xz	1 L'	1 g	Apply as a spot spray to all areas
institutions and food handling establishments	?÷,	1( ,	1 q'	î q	frequented by reckmaches
Restuarants Supermarkets Or mist spray inside buildings for German conkroaches	•-	K 12	' p'	1 q	Difute with 34 and apply by remote control when buildings are not occupied. Cover exposed food and equipment Use togal diduted spray per 50,000 cu. ft.
Food Processing Plants	45,	4 12	1 p1	¹ q	Spray when not in liperat in Civer
	7.3	10 2	t gt	2 g:	priti e ard equipment
Meat Packing Plants	2°,	.u s	1 21	∂ ធ¹	Spray order and nedship product
Poultry Processing Plants	? -	11 -2	• 4•	3 d.	Spray later and medible product areas
MISCELLANEOUS INSECTS Fleas and spiders in buildings laundry trucks dog kennels etc.	••	5 7	1 p.	1 g	Apply tight fiduated spray per 1000 sq ft. Spray flours, walls cracks crevices sleeping quarters of ets where inserts bried and huf-bor.
Bedbugs in buildings and sleeping quarters on beds and biolding	14.	52	1 pt	1 01	Spray mattress and bedding lightly countries and folding cracks and folding tracks and residence in bed frames walls baselinards and floors.
Figas in lawn and kennel runs	1°.	57	1 p'	t d:	Spray infested areas. Keep children off-treated area with spray has dried
Ants Crickets Mosquitoes and	1^2	5 02	1 pt	f q!	Spray areas where insects frequent

Ants, Grickets, Mosquitoes, and

CODE NO. 18-1101

U.S.D.A. Reg. No. 1990-328



For flies, lice, and pest control use. Water emulsifiable and oil soluble.

Active Ingredients: Ronnel (0,0-dimethyl 0-(2,45-Trichlorophenyl) Phosphorothioate +)	24%
Aromatic Petroleum Derivative Solvent	61%
Inert Ingredients	
*Korlan (i) Trademark of the Dow Chemical Company	00 /0

# CAUTION

· From technical grade. Not suitable for internal use in animals

Keep out of reach of children
See side panel for additional precautionary labeling

**NET CONTENTS 5 GALLONS** 

#### CAUTION

Injurious to the eyes. Irritating to the skin. Harmful if swallowed. Keep out of reach of children

Do not get in eyes. Avoid prolonged or repeated contact with the skin. Avoid prolonged breathing of spray mists Avoid contamination of food, feedstuffs, and water supplies.

Keep material out of reach of animals. Keep away from heat and open flame. Keep container closed. Wash thoroughly with soap and water after handling.

### COMBUSTIBLE LIQUID

Note to physician: This material is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

#### WARNING

Do not spray hogs within 6 weeks and sheep and goats within 12 weeks of slaughter. Do not spray beef cattle within 8 weeks of slaughter. Do not spray lactating dairy animals, or dry dairy animals within 21 days of freshening. Back rubbers may be used for beef animals to within 2 weeks of slaughter, or dry dairy cows to within 3 weeks of freshening. Not for use on lactating dairy cows. Do not allow spray to come in contact with food, feed or water supplies. Remove lactating dairy cows from buildings while applying residual fly sprays. Do not use on animals receiving an organic phosphate treatment from any other source at the time of treatment.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

## **KORLAN 24E Livestock Remedy**

## - Not For Human Use

Notice: Seller warrants that the product conform to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

U. S Patent No. 2,599,516.

