POA ANNUA: To control annual bluegrass and other erratic or late germinating grasses and weeds, a late summer or early fall application of 4-6 pounds per acre of CO-OP Lawn and Garden Weed Killer should supplement early spring application.

The use of CO-OP Lawn and Garden Weed Killer on bentgrass should be restricted to portions of golf courses infested with crabgrass and other annual grasses and weeds. Do not apply CO-OP Lawn and Garden Weed Killer to bentgrass or fairway turf. Repeated use is not recommended. At the recommended rate of application, CO-OP Lawn and Garden Weed Killer does not show herbicidal activity in soil. Normally it is safe to plant seeds or transplants no more than the recommended rate of the weed.

FLOWER GARDENS

Apply at transplant, or as supplemental application at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply to the soil for annual establishment or selected out of flower beds. Care should be taken not to apply in the spring before all seedlings have emerged. Beds should be cultivated at any time prior to this. If beds have been cultivated and are free of weeds, weed control up to four months or more may be expected following proper application.

NURSERY STOCK OR ORNAMENTAL PLANTINGS

Apply at transplant, late fall, early spring at the rate of 180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Applications should be made to soil recently cultivated to a uniform texture. These can be made prior to following transplanting. With established plantings, applications should be made following proper cultivation to remove existing weeds. Where possible, this should be done early in the spring. Late summer applications may provide control of fall germinating weeds if made following cultivation. Weed control of three months or more may be expected following proper application. See label.

VEGETABLES

Broccoli, Brussels sprouts, cauliflower and cabbage: Apply at transplanting or layby at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply uniformly to the soil. CO-OP Lawn and Garden Weed Killer can be applied directly over transplants without injury. If weeds have emerged, crops should be clean cultivated or weeded prior to application. **Beans (dry and snap). Southern Peas (blackeyed):** Apply at seeding or transplanting at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply uniformly to the soil at time of seeding.

Cucumbers, squash (winter and summer), melons (cantaloupe, watermelon, honeydew): Apply 1-3 weeks after seeding at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply uniformly to the soil 4 to 6 weeks after seeding. Crops should be cultivated or weeded prior to the application if weeds have emerged.

Collards, kale, mustard greens, turnips (greens and roots): Apply at seeding at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply uniformly to the soil at time of seeding. **Garlic and onions:** Applications at seeding or transplanting or layby applications can be made as crops up to 14 weeks after seeding or transplanting at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply uniformly to soil. CO-OP Lawn and Garden Weed Killer can be applied directly over transplants without injury. Layby applications can be made and may up to 14 weeks after planting of transplants. Should weeds emerge prior to layby, crops should be cultivated or weeded prior to application.

Lettuce (leaf): Apply 1-3 weeks after seeding at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply uniformly to the soil. Apply not later than three weeks after emergence.

Lettuce (head): Apply 1-3 weeks after seeding at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply only to soil. Apply only between 1 to 6 weeks after seeding.

Potatoes (whole or cut pieces): Apply at planting, dropoff or layby at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply uniformly to the soil at planting. For layby applications, the soil should be cultivated and free of weeds. CO-OP Lawn and Garden Weed Killer should be applied after crop is established. Layby applications can be made up to nine weeks after planting. If weeds have emerged at layby, the crop should be weeded or cultivated prior to application.

Sweet potatoes and yams: Apply at transplanting or layby at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply uniformly to the soil at transplanting. CO-OP Lawn and Garden Weed Killer can be applied directly over transplants without injury. Layby applications can be made up to 6 weeks after transplanting. If weeds have emerged, the crop should be cultivated or weeded prior to making the CO-OP Lawn and Garden Weed Killer application.

Tomatoes: Apply 4-6 weeks after transplanting at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Application should be confined to a period of 4 to 6 weeks after transplanting. CO-OP Lawn and Garden Weed Killer can be applied directly over the transplants without injury. If weeds have emerged, the crop should be cultivated or weeded prior to making the CO-OP Lawn and Garden Weed Killer application.

Peppers and peppers: Apply at transplanting or layby at the rate of 160-180 pounds CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply uniformly to the soil at transplanting. CO-OP Lawn and Garden Weed Killer can be applied directly over transplants without injury. Layby applications can be made up to 6 weeks after planting. If weeds are present, the crop should be cultivated or weeded prior to making the layby application.

Strawberries (new plantings): Apply at planting at the rate of 180 lbs. CO-OP Lawn and Garden Weed Killer per acre (For established plantings): Apply in fall and early spring at the rate of 160-180 lbs. CO-OP Lawn and Garden Weed Killer per acre (4 pounds per 1,000 square feet). Apply uniformly to the soil. For new beds apply 180 pounds at time planting. Applications to established plantings should be made in fall and early spring. Applications of 160-180 pounds per acre may be applied to control late summer or early fall germinating weeds. DO NOT apply after first bloom. These applications can be made directly over plants without injury.

CO-OP Lawn and Garden Weed Killer is specifically recommended for weed control with the following:

Alpha, Arborvitae, Ash, Azalea, Barberry, Boxwood, Birch, Camellia, Chestnut, Coleosader, Cottonwood, Crab Apple, Cypress, Deutzia, Dockand, Elaeagnus, Elm, Eucalyptus, Fir, Forsythia, Gum, Hawthorn, Holly, Hydrangea, Jap. anese, Juniper, Larkspur, Lilac, Locust, Magnolia, Maple, Mockorange, Mountain, Mulberry, Oak, Pachysandra, Pine, Pittosporum, Poplar, Poplar, Privet, Redbud, Rhododendron, Rose, Russian olive, Spirea, Spruce, Sycamore, Tree peony, Tulip tree, Varnish tree, Viburnum, Walnut, Weigela, Willow, Yew, Bellflower, Bentgrass, Bindweed, Honeysuckle, Pachysandra, Wintercreeper, Ageratum, Alyssum, Aster, Bittercress, Bluebell, Cardinal flower, Chickweed, Bluegrass, Canada thistle, Candytuft, Cinquefoil, Chrysanthemum, Columbine, Coleus, Coreopsis, Cornflower, Creeping Charlie, Dandelion, Daylily, Delphinium, Dianthus, Dichondra, Dock, Dusty miller, Fleabane, Forget-me-not, Foxglove, Garden spurge, Geranium, Gerbera, Gladiolus, Goldenrod, Gypsophila, Hawkweed, Hollyhock, Iris, Lambsquarters, Lantana, Lavender, Morning glory, Mullein, Nutsedge, Oxalis, Pansy, Pennywort, Periwinkle, Phlox, Plantain, Poppy, Purslane, Quackgrass, Ragweed, Ragwort, Shepherdspurse, Smartweed, Speedwell, Spurge, Stinging nettle, Sunflower, Sweet William, Thistle, Violet, Wild buckwheat, Wild garlic, Wild mustard, Yarrow, Yellow rocket.

DO NOT USE on lawns, flowers, ornamentals or vegetables unless listed on this label. Do not use on lawns, flowers or other plantings that are not listed on this label. Do not apply under hot dry conditions or to stressed turf because injury may result. Do not use on newly seeded lawns until grass has been mowed at least three times. Do not apply to lawns for 30 days following application of 2,4-D type herbicides. Do not apply after July 15.

CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse. Keep out of reach of children. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply under conditions or in a manner that will result in runoff or drift to non-target areas. Do not apply when weather conditions favor runoff or drift. Do not allow people or pets on treated turf until sprays have dried. Do not feed treated grass clippings to livestock. Do not apply more than 20 lbs. product per 1,000 square feet per application. Do not apply more than 2 applications per year.

WARRANTY AND LIMITATION OF DAMAGES

SELLER warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. SELLER makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY, and no agent of SELLER is authorized to do so except in writing with specific reference to this warranty. In no event shall SELLER'S liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

Manufactured by

FARMLAND INDUSTRIES, INC.

Kansas City, Missouri