

Renovation of existing stand of big bluestem and switchgrass planted on CRP acres.

Broadcast 1-2 qt. / A to existing stands of big bluestem and switchgrass prior to the emergence of weeds. Use the low rate on soils containing from 1-2% organic matter. Use the high rate on soils with 2% or greater organic matter content.

Renovation of existing stands of the following perennial range grasses planted on CRP acres: Blue grama, Indiangrass, little bluestem, sand lovegrass, sideoats grama and western wheatgrass.

Broadcast 0.5 - 1 qt / A in the spring prior to weed emergence, or in the fall before the ground freezes and prior to weed emergence after these species have been established for at least 1 growing season, for control or partial control of the weeds listed above. Use the low rate for weeds that are controlled or suppressed easily.

Aerial Application: Make applications at a maximum height of 10 ft above vegetation. Use low drift nozzles at a maximum pressure of 40 psi. To control drift, restrict application to periods when wind speed does not exceed 10 mph. To assure that drift will not adversely affect adjacent sensitive nontarget plants, apply Liquid Atrazine 4L by aircraft at a minimum upwind distance of 400 ft. from sensitive plants. Use 3-5 gallons / Acre total water volume when a dense heavy ground cover is present.

Note: (1) Do not cut or feed grass hay to livestock. (2) Do not graze treated areas. (3) Do not use seeds for bird food. (4) Do not dump or spill product or dispose of product containers within the reach of livestock (5) Follow all applicable restrictions of the Conservation Reserve Program.

Precautions: To avoid crop injury, make only one (1) application per year. Slight discoloration of desirable grasses may occur following treatment. Injury may be enhanced when used on neutral or alkaline soils.

SEVERE DROUGHT CONDITIONS

Do not graze forage or cut forage for hay from CRP acres if treated with atrazine. Under severe drought conditions, the Conservation Reserve Program allows grazing and making of hay from CRP acres, as so specified by the local Agricultural Stabilization & Conservation Service (ASCS) office. This label does not allow for grazing or making of hay from atrazine treated acres, under those or any other circumstance.

SUGARCANE

For control of many broadleaf weeds and grass weeds, including amaranths, crabgrass, fireweed, Flora's paintbrush, foxtails, junglerice and wiregrass, apply 4-5 pts. / A of Liquid Atrazine 4L at time of planting or ratooning, but before sugarcane emerges. For aerial application, broadcast in a minimum of 5 gal / A spray volume. For application with by ground, broadcast or band, apply in a minimum of 20 gal / A, unless indicated otherwise. One (1) additional application may be made over the sugarcane as it emerges, and two (2) additional applications may be made interline, after emergence, as directed sprays. Repeat treatments, where needed, may be applied broadcast, band, or interline as suggested with the final application being made prior to "close-in".

Do not apply more than 10 quarts / A to any one crop of sugarcane.

NOTE: Where high rates of Liquid Atrazine 4L are used alone, apply in a minimum of 1 qt of water for each 1 qt of Liquid Atrazine 4L applied per acre.

Aerial Application: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft., using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speeds do not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply Liquid Atrazine 4L alone by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

Florida

For control of emerged pellitory (artillery) weed: Apply 0.8 - 1.2 pints / A in at least 40 gallons of water as a directed spray, by ground equipment, prior to "close-in". Add 4 quarts of surfactant for each 100 gallons of spray. Thoroughly wet weed foliage with spray.

For control of alexandergrass, large crabgrass, pellitory (artillery) weed, and spiny amaranth, use one of the following methods at planting or ratooning:

1. Apply 8 pts. / A preemergence. Follow with 1 or 2 applications, as needed, postemergence to sugarcane and weeds, at 4 pts. / A. Treat before weeds exceed 1.5 inches in height.
2. Apply 1-3 times, as needed, at 4 pts. / A postemergence to sugarcane and weeds. Treat before weeds exceed 1.5 inches in height.

Louisiana

For control of annual weeds during the summer fallow period, apply 4 pts. / A to weed-free beds immediately after bed formation. Follow normal weed control program after planting.

Precautions: If making 4 pts. / A applications during the summer fallow period, do not exceed 8 qt. / A during the remainder of the season in order to avoid crop injury and/or illegal residues.

Texas

Use Liquid Atrazine 4L for control of barnyardgrass, pigweed, purslane, and sunflower, in plant or ratoon sugarcane.

Apply 8 pts. / A of Liquid Atrazine 4L 4L preemergence. Follow with 1 or 2 applications as needed, at 6 pts. / a postemergence to sugarcane and weeds.

For best results, when weeds are emerged, add a nonionic surfactant at a concentration of 2 qts. / 100 gals. to the spray and apply before weeds exceed 1.5 inches in height.

Precautions: (1) Injury to sugarcane may occur when under moisture stress, when soil is of low adsorptive capacity, or when land is first cropped to sugarcane. (2) Do not apply after close-in.

TURFGRASS FOR SOD (Florida Only)

St. Augustinegrass, Centipedegrass, and Zoysiagrass

Broadcast 4-8 pts. / A according to soil texture (Table 4.) to control those weeds listed under "General Information" for Liquid Atrazine 4L applied alone.

) Table 4. Broadcast Rates of Liquid Atrazine 4L for use on Turfgrass in Florida

Muck or Peat	8 pints	Old beds: Within 2 days after lifting sod.
		New beds: 3-4 days after sprigging or plugging
Sandy Soil	4 pints	Old beds: Within 2 days after lifting sod.
		New beds: 7-10 days after sprigging or plugging.

If weeds regrow, apply an additional 4 pts. / A on muck or peat, or 2 pts./ A on sandy soil.

Precautions: To avoid crop injury, (1) Do not apply within 30 days prior to cutting or lifting. (2) Do not apply in combination with surfactants or other spray additives. (3) Use only on turfgrass reasonably free of infestations of insects, nematodes and diseases. (4) On newly sprigged turfgrass, temporary slowing of growth may follow applications.

TURFGRASS FOR FAIRWAYS, LAWNS, SOD PRODUCTION (in states other than Florida*) AND SIMILAR AREAS.

• For use on turfgrass for sod in Florida, see "TURFGRASS FOR SOD (Florida Only)" in above section.

Bermudagrass, Centipedegrass, St. Augustinegrass and Zoysiagrass

Apply Liquid Atrazine 4L after October 1, before emergence of winter annual weeds, for control of annual bluegrass, burclover, carpet burweed, chickweed, corn speedwell, henbit, hop clover and sr rweed. Liquid Atrazine 4L will control annual bluegrass even if it is emerged at the time of treatment. For control of summer annual weeds listed "GENERAL INFORMATION" section of this label, also apply Liquid Atrazine 4L in late winter before the weeds emerge. Apply in a minimum of 15 gallons of water per acre or 1 gallon per 1,000 square feet.

Where annual bluegrass is the major weed, use 1 qt. / A at least 30 days after the previous application, but not after April 15. However, do not make more than two (2) applications of this herbicide per year.

Precautions: On newly sprigged turfgrass and hybrid bermudagrass, temporary slowing of growth and yellowing may occur following application. To avoid turf injury,

- (1) Use only on turfgrass reasonably free of infestations of insects, nematodes and diseases.
- (2) Do not use on golf greens.
- (3) Do not use north of North Carolina, (with the exception of the Virginia Coastal Plain) or west of the high rainfall areas of Eastern Oklahoma and Eastern Texas.
- (4) Do not use on muck or alkaline soils.
- (5) Do not apply over the rooting area of trees or ornamentals not listed on this label.
- (6) Do not overseed with desirable turfgrass within 4 months before or 6 months after treatment.
- (7) Do not apply this product to newly seeded Bermudagrass until it has overwintered and has a well-developed rhizome system. Do not exceed 2 qts. / A of this product within 12 months of seeding Bermudagrass.

Note: Do not graze or feed turf clippings to animals, or illegal residues may result.

MACADAMIA NUTS

For preemergence control of many broadleaf and grass weeds, including crabgrass, foxtail, wiregrass, Florida's paintbrush, spanishneedles and fireweed, broadcast 4-8 pts. / A of Liquid Atrazine 4 L before harvest and before weeds emerge. Repeat as necessary. Do not spray when nuts are on the ground or during the harvest period. Do not apply by air.

GUAVA

) Use only on established plantings which are at least 18 months old. Apply as a directed spray, at 4-8 pts. / A of Liquid Atrazine 4L in 20-50 gallons of spray mix, preemergence or early postemergence to weeds. When applying postemergence, the use of a surfactant and greater spray volume (80-100 gallons of spray mix per acre) may enhance weed control. This product controls many annual broadleaf and grass weeds, including fireweed, purslane, scarlet pimpernel, spanishneedles and sowthistles.

Note: To avoid illegal residues, (1) Do not allow spray to contact guava foliage or fruit. (2) Do not apply more frequently than at 4 month intervals. (3) Do not apply more than 16 pints of Liquid Atrazine 4L per acre per year.

CONIFERS

For control of annual broadleaf and grass weeds prior to transplanting, after transplanting, or in established conifers [including Douglas fir, grand fir, noble fir, white fir, Austrian pine, bishop pine, Jeffrey pine, knobcone pine, loblolly pine, lodgepole pine, (shore pine), monterey pine, ponderosa pine, Scotch pine, slash pine, blue spruce and Sitka spruce]: Broadcast 4-8 pints in a minimum of 5 gallons of water per acre by air or 10 gallons of water per acre by ground, before weeds are 1.5 inches tall. Apply to established trees between fall and early spring while trees are dormant. For new transplants, apply during or soon after transplanting. For applications prior to transplanting wait for sufficient precipitation to activate Liquid Atrazine 4L before transplanting. In areas where spring and summer rainfall is inadequate to activate Liquid Atrazine 4L, apply during fall prior to spring transplanting.

For the list of weeds controlled, see "GENERAL INFORMATION" section of this label.

8/29/94

Quackgrass control: Broadcast 8 pints in a minimum of 5 gallons of water per acre by air or 10 gallons per acre by ground, between fall and early spring while trees are dormant and before quackgrass is more than 1.5 inches in height.

Precautions:

- (1) In areas west of the Rocky Mountains (except the Great Basin) grazing may begin 7 months after a fall application, or 3 months after a winter or spring application.
- (2) To prevent illegal residues, do not graze treated areas of the Great Basin or east of the Rocky Mountains.
- (3) Temporary injury to trees may occur following use of Liquid Atrazine 4L on coarse textured soil.
- (4) To avoid crop injury, do not apply to seedbeds.
- (5) Apply only once per year.

Aerial Application: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft., using low drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speeds do not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply Liquid Atrazine 4L alone by aircraft at a minimum upwind distance of 400 ft. from sensitive plants.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried

ROADSIDES

Apply Liquid Atrazine 4L control certain annual weeds in established perennial grasses along roadsides in CO, KS, MT, ND, NE, SD and W., including cheatgrass (Downy brome, chess) common (annual) broomweed, little barley, medusahead, sagewort, and tumble mustard. Broadcast this product at a rate of 2 pts. / A in a minimum of 10 gallons of water by ground equipment, in the fall, before the ground freezes, or after thawing in the spring before the established grasses green-up and before weeds emerge. Examples of established desirable grasses include big bluestem, bluegrama, bromegrass, buffalograss, crested wheatgrass, Indiangrass, little bluestem, side-oats grama, switchgrass and Western wheatgrass. Apply only once per year. Temporary discoloration or other form of injury to the desirable perennial grasses may occur following application.

NOTE: Do not cut or feed road-side hay. Do not allow livestock to graze in treated areas.

Caliber 90, Concep, Duale, and Prince are trademarks of Ciba-Geigy Corporation
Compex is a trademark of KALO Agricultural Chemical, Inc.
Gramoxone and Sutan are trademarks of Zeneca Ag Products
Lasso, Ramrod and Roundup are trademarks of Monsanto Company
Unite is a trademark of HACO, Inc.
X-77 is a trademark of Chevron Chemical Company

Reg # 1990-381

PM 25

1920

**RESTRICTED USE PESTICIDE
(GROUND AND SURFACE WATER CONCERNS)**

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATIONS.

THIS PRODUCT IS A RESTRICTED USE HERBICIDE DUE TO GROUND AND SURFACE WATER CONCERNS. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE TO REACH GROUND AND SURFACE WATER.

EPA Reg. No. 1990-381
EPA Est. No. 32761-MO-3

FARMLAND Liquid Atrazine 4L

Herbicide - for season-long weed control in corn, sorghum, and certain other crops

CAUTION
KEEP OUT OF REACH OF CHILDREN
For Professional or Agricultural Use Only

Shake well before using.

Directions for use
See inside of booklet for complete directions for use and additional precautionary statements.


Net Contents: 2.5 Gallons
U.S. Standard Measure
Use entire contents at one time.

Manufactured by Farmland Industries, Inc.
General Office Kansas City, MO 64116

ACTIVE INGREDIENTS:

Atrazine, 2-chloro-4-ethylamino-6-isopropylamino-s-triazine	40.8%
Related compounds	2.2%
INERT INGREDIENTS	57.0%
TOTAL	100.0%


Liquid Atrazine 4L contains 4 lbs. active ingredients per gallon.



Revision Date 8/94

8/29/94

2920



PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Caution: Keep out of reach of children

Harmful if swallowed. Do not get in eyes. Avoid contact with skin, inhalation of vapors or spray mist and contamination of food and feed.

Personal Protective Equipment (PPE):

Handlers who may be exposed to the dilute through application or other tasks must wear:

- A. Long-sleeved shirt and long pants
- B. Waterproof gloves
- C. Chemical-resistant footwear plus socks

Handlers who may be exposed to the concentrate through mixing, loading, application or other tasks must wear:

- A. long sleeved shirt and long pants
- B. Waterproof gloves
- C. Chemical-resistant footwear plus socks
- D. Protective eyewear
- E. Chemical-resistant apron when mixing or loading.

Follow Manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- * Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- * Remove clothing immediately if pesticide gets inside. then wash thoroughly and put on clean clothing.
- * Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Statement of Practical Treatment

First Aid Procedures: If poisoning is suspected, do not wait for symptoms to develop. Immediately begin the recommended procedures below. Contact a physician, the nearest hospital or the nearest Poison Control Center. Tell the person contacted the type and amount of exposure, describe the symptoms and follow the advice given.

Note to Physician: There is no specific antidote for atrazine. If this product is ingested, induce emesis or lavage stomach. The use of an aqueous slurry of activated charcoal may be considered.

If Swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger, or if available, by administering Syrup of Ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

8/29/94

On Skin: Wipe away excess material from the skin. Wash with plenty of soap and water. Do not attempt to neutralize with chemical agents. Get medical attention.

In Eyes: Immediately flush the eyes with large amounts of running water for a minimum of 15 minutes. Hold the eyelids apart during the flushing to ensure rinsing of the entire surface of the eyes and lids with water. Do not attempt to neutralize with chemical agents. Obtain medical attention if irritation occurs.

If Inhaled: Remove victim to uncontaminated atmosphere. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

Environmental Hazards

Groundwater Protection: Atrazine can travel (seep or leach) through soil and can enter groundwater which may be used as drinking water. Atrazine has been found in groundwater. Users are advised not to apply Atrazine to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater. Groundwater contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

) This product may not be mixed/loaded, or used within 50 feet of all wells including abandoned wells, drainage wells, and sink holes.

Operations that involve mixing, loading, rinsing or washing of this product into or from pesticide handling application equipment or containers within 50 feet of any well are prohibited unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spill or equipment leaks, container or equipment rinse or wash water and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means that the pad must be self self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum of 100% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad, shall have a minimum containment capacity of 100% of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing or loading sites.

States may have in effect additional requirements regarding wellhead setbacks and operational area containment.

) This product may not be mixed or loaded within 50 feet of intermittent streams and rivers, natural or impounded lakes and reservoirs. This product may not be applied aerially or by ground within 66 feet of the points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 ft. around natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or setback from runoff points must be planted to crop or seeded with grass or other suitable crop.

Do not apply this product through any type of irrigation system.

Fish and Wildlife Hazards: This pesticide is toxic to aquatic invertebrates. For terrestrial uses, do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash waters.

Directions For Use

It is a violation of FEDERAL LAW to use this product in a manner inconsistent with its labeling.

Important: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during applications. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENT

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- a. Coveralls
- b. Waterproof gloves
- c. Chemical resistant footwear plus socks

Storage and Disposal

Storage

Keep This Material Out of the Reach of Children

Store only in the original container with cap securely tightened.

Containers must be stored in a cool, dry, well-ventilated area. Store away from food, feeds and fertilizers. Do not contaminate water, food, feed by storage or disposal.

Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

Spill or Leak Procedures: In the event of a spill or leak, make sure all personnel involved in spill cleanup follow good industrial hygiene practices. Small spills can be handled routinely. Cover the spill with an absorbent material such as vermiculite or sawdust to prevent dust. Sweep up the material and place in an appropriate chemical waste container. Seal container and dispose of according to "Pesticide" disposal instructions below. Wash the spill area with a saturated solution of sodium carbonate and a strong detergent. Flush the spill area with water to remove any residue. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-888-8372 day or night.

Disposal

Containers: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state authorities, by burning. If burned, stay out of smoke.

Pesticide: Do not contaminate water, food, or feed by storage or disposal, or cleaning of equipment. Open dumping is prohibited. Wastes resulting from the use of this product are acutely toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

Conditions of Sale and Warranty

Conditions of Sale. This product is sold subject to and buyer and users are deemed to have accepted the terms contained in the CONDITIONS OF SALE AND WARRANTY, DIRECTIONS FOR USE AND PRECAUTIONARY

5720

STATEMENTS, which may be varied only by agreement in writing by a duly authorized representative of FARMLAND INDUSTRIES, INC.

Inherent Risks: The directions for use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as presence of other materials or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of FARMLAND INDUSTRIES, INC., or the seller. All such risks shall be assumed by the buyer.

Warranty Limitations: FARMLAND INDUSTRIES, INC., warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the DIRECTIONS FOR USE, subject to the inherent risks referred to above. FARMLAND INDUSTRIES, INC., MAKES NO OTHER EXPRESSED OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESSED OR IMPLIED WARRANTY.

Limitation of Liability: In no case shall FARMLAND INDUSTRIES, INC., or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product. No claim of any kind shall be greater in amount than the purchase price of the product in respect of which such damages are claimed.

Do not reuse empty container. Apply this product only as specified on this label.

General Information

LIQUID ATRAZINE 4L herbicide will control many annual broadleaf and grass weeds in corn, sorghum, sugarcane and certain other crops specified on this label.

Broadleaf and Grass Weeds Controlled with Liquid Atrazine 4L Alone - Corn, Grain Sorghum and other crops specified on this label.*

Preplant Surface-Applied, Preplant Incorporated, or Preemergence (or Postemergence at 4 pts. / A with Oil)

Broadleaf and Grass Weeds Controlled:

Annual Morningglory	Barnyardgrass (watergrass)***	Cocklebur**
Giant Foxtail**	Green Foxtail***	Groundcherry
Kochia	Lambsquarters	Large (hairy) Crabgrass**
Mustards	Nightshade	Pigweed
Purslane	Ragweed	Sicklepod**
Velvetleaf**	Wild Oats	Witchgrass (Panicum Capillare)***
Yellow Foxtail***		

Postemergence with Emulsifiable Oil or Oil Concentrate in Water (at 2.4 pts. / A)

Broadleaf Weeds Controlled:

Annual Morningglory	Cocklebur	Jimson Weed
Lambsquarters	Mustards	Pigweed
Ragweed	Smartweed	Wild buckwheat
Velvetleaf**		

*Where there are state/local requirements regarding atrazine use (including lower maximum rates and/or higher set-backs) which are different from the label, the more restrictive/protective requirements apply. Certain states may have established rate limitations within specific geographical areas.

Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

** Partial control only

*** Partial control only on medium- and fine-textured soils.

Atrazine may be applied before or after weeds emerge. The continuous use of this product and other triazines may cause the development of biotypes of grasses and broadleaf weeds that cannot be effectively controlled by this and

8/29/94

related herbicides. Where this is known or suspected, and weeds controlled by this product are expected to be present along with resistant biotypes, LIQUID ATRAZINE 4L should be either tank mixed with or used in sequence with a herbicide approved for use with Atrazine which is not in the triazine class of chemicals. If only resistant biotypes are expected to be present use a registered non-triazine herbicide. Read and follow all directions for use and precautions on the labels of products to be tank mixed with Atrazine. Consult your State Extension Service for specific recommendations for your area.

In each case where a range of rates is given, the lower rate should be used on coarse-textured soils and soils low in organic matter, and the higher rate should be used on fine-textured soils and soils high in organic matter. Since Atrazine acts mainly through root absorption, its effectiveness depends on rainfall or irrigation to move it into the root zone. Should weeds develop, a shallow cultivation or rotary hoeing will generally result in better weed control.

This product is nonflammable, noncorrosive to equipment and metal surfaces, and has low electrical conductivity.

Do not apply this product through any type of irrigation system.

Care should be taken to avoid using Atrazine where adjacent desirable trees, shrubs or plants may be injured. Avoid using near adjacent desirable plants or in greenhouses, or injury may occur.

NOTE: FARMLAND INDUSTRIES, INC., does not recommend the use of LIQUID ATRAZINE 4L in combination with other herbicides or oils, except as specifically described on this label.

APPLICATION PROCEDURES

Ground Application: Use conventional ground sprayers equipped with nozzles that provide accurate and uniform application. For the most uniform distribution of broadcast applications, use 80 degrees flat fan-type nozzles. For band applications, use flat fan even-spray nozzles. Be certain that nozzles are uniformly spaced and are of the same size and type. Calibrate sprayer before use and recalibrate at the start of each season and when changing carriers. Screens and strainers should be no finer than 50 mesh between the pump and bowl. Screens on the suction side of the pump should be 15 mesh or coarser. Do not place a screen in the recirculation line.

Use a pump with capacity to: (1) provide sufficient hydraulic agitation during mixing and application to keep the material in suspension (2) maintain 30-40 psi operating pressure, and (3) provide a minimum of 20% by-pass at all times. (At minimum, 10 gallons / minute per 100 gallon tanks size). Unless otherwise specified, use a minimum of 5 to 10 gallons of water per acre for preplant incorporated, pre-emergence and post (with or without oil or surfactant) applications.

For band applications, calculate the amount to be applied per acre as follows:

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Rate/A for Broadcast Treatment} = \text{Amount needed for Band Treatment}$$

Aerial Application: For preplant and pre-emergence broadcast treatments, apply at a 1:1 ratio LIQUID ATRAZINE 4L to water (example: recommendation calls for 1 quart of product and is mixed with 1 quart of water) to be applied per acre. For postemergence treatments on corn and sorghum, apply the recommended rate of LIQUID ATRAZINE 4L plus oil in a minimum of 2 gallons of water per acre.

Avoid applications under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. The above water volumes are the minimum to be applied per acre.

LIQUID ATRAZINE 4L in Water Applications: LIQUID ATRAZINE 4L, a liquefied formulation, should be mixed with water and applied as a spray. Pour LIQUID ATRAZINE 4L into the tank during or after filling. Hydraulic (jet) or mechanical agitation is recommended during mixing and application to keep the material in suspension. All return lines to the tank must discharge below liquid level and agitation should not be so violent as to cause air bubbles to form in the liquid. Wash sprayer thoroughly after use.

LIQUID ATRAZINE 4L in Liquid Fertilizer Application: Nitrogen solutions or complete liquid fertilizers may replace all or part of the water as a carrier for pre-emergence, preplant incorporated, or preplant surface ground application on corn and sorghum. Mixing should be accomplished as described under "Water Applications". Do not apply in nitrogen solution or complete liquid fertilizer after corn or sorghum emerges or crop injury may occur. Check the compatibility of this product with liquid fertilizer and/or nitrogen solution, as shown below, before use.

Compatibility Test: Always check compatibility of liquid fertilizers with herbicide(s) each time before use. Be especially careful when using complete suspension or fluid fertilizers as serious compatibility problems are more likely to occur. Commercial application equipment may improve compatibility in some instances. The following test assumes a spray volume of 25 gal / A. For other spray volumes, make appropriate changes in the ingredients.

Check compatibility using this procedure:

1. Add 1 pt. of fertilizer to each of two (2) 1- qt. jars with tight lids.
2. To one of the jars, add 1/4 tsp. or 1.2 milliliters of a compatibility agent such as Compex or Unite (1/4 tsp. in 1 pt. is equivalent to 2 pt. per 100 gal of spray). Shake or stir gently to mix.
3. To both jars add the appropriate amount of herbicide (see below). If more than one herbicide is used, add them separately (dry herbicides first, flowables next and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix. The appropriate amount of herbicide for this test is:
Dry herbicides - For each 1 lb per acre rate, add 1.5 level tsp to each jar.
Liquid Herbicides - For each 1 pt. per acre rate, add 0.5 tsp (2.5 milliliters) to each jar.
4. After adding oil ingredients, put lids on tightly and invert each jar 10 times to mix. Let the mix tures stand 15 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, try adding half the compatibility agent to the fertilizer and half to the herbicide, before combining the herbicide and fertilizer. If incompatibility is still observed, do not use the mixture.

LIQUID ATRAZINE 4L Applications in Water Plus Emulsifiable Oil or Oil Concentrate: The addition of emulsifiable oil (petroleum-derived, petroleum-derived oil concentrate, or single or mixed crop derived oil concentrate) to post emergence Atrazine in water sprays may speed the activity of Atrazine and provide quicker kill of weeds. However, under certain conditions, the use of either type of oil may seriously damage corn or sorghum. To minimize the possibility of damage, follow the directions, procedures and precaution given below.

Use one of the following properly emulsified:
A suitable oil concentrate containing at least 1%, but not more than 20%, suitable emulsifier or surfactant blend.
A petroleum derived oil containing at least 1% suitable emulsifier.

Note: In the event of a compatibility problem when mixing oil with this product and water, a compatibility agent such as Compex or Unite should be used. Any of the above oils contaminated with water or other materials can cause compatibility problems and/or crop injury.

Mixing Procedures - all uses:

- (1) Thoroughly clean sprayer prior to use. Do not use sprayer contaminated with 2, 4-D or other materials as crop damage or sprayer clogging may result.
- (2) Fill spray tank one-fourth full with clean water, nitrogen solution or complete liquid fertilizer .
- (3) With the agitation system in operation, such that there is a rippling or rolling action on the liquid surface, add the Atrazine. Continue filling tank with water, nitrogen solution, or complete liquid fertilizer, until 90% full. Increase agitation if necessary to maintain surface action
- (4) Add emulsifiable oil, oil concentrate, or tank mix herbicides last, after this product is thoroughly suspended. Finish filling tank

- (5) Empty the tank as completely as possible before refilling to prevent a buildup of oil or emulsifiable concentrate residue in the tank. Keep the agitation system in operation to avoid separation of materials in any mixture remaining in the tank.
- (6) If an oil or emulsifiable concentrate film starts to build up in the tank, drain the tank and clean out with a strong detergent solution or solvent.
- (7) Clean the sprayer thoroughly immediately after using by flushing the system with water containing a detergent.

Do not apply this product through any type of irrigation system.

Rotational Crop Suggestions - all uses:

- (1) Land treated with Liquid Atrazine 4L should not be planted to any crop except corn or sorghum until the following year or injury may occur.
- (2) If Atrazine is applied after June 10, do not rotate with crops other than corn or sorghum the next year or crop injury may occur.
- (3) In the high plains or intermountain areas of the west where rainfall is sparse and erratic or where irrigation is required, use Liquid Atrazine 4L only when corn or sorghum is to follow corn or sorghum, or a crop of untreated corn or sorghum is to precede other rotational crops.
- (4) In western Minnesota and eastern parts of the Dakotas, Nebraska and Kansas, corn or sorghum treated with Liquid Atrazine 4L should not be followed with soybeans if the broadcast rate applied was more than 4 pints per acre (or comparable rate in a band) or soybean injury may occur.
- (5) Injury may occur in soybeans planted the year following a Liquid Atrazine 4L application on soils having a calcareous surface layer.
- (6) Do not plant sugarbeets, tobacco, vegetables (including dry beans), spring-seeded small grains, or small-seeded legumes and grasses the year following Liquid Atrazine 4L application or injury may occur.

DIRECTIONS FOR USE

CORN:

LIQUID ATRAZINE 4L ALONE Apply either before planting, at planting or after planting as indicated below. For soil applications prior to crop emergence (i.e., early preplant, preplant incorporated, preplant surface, at planting or preemergence) see table 1.

Preplant Surface-Applied (Broadleaf and grass control): Use on medium- and fine-textured soil with minimum-tillage or no tillage systems only in CO, IL, IN, IA, KS, KY, MN, MO, MT, NE, ND, SD, WI, and WY. Apply the recommended rate of Liquid Atrazine 4L shown in Table 1 up to 45 days prior to planting. On coarse textured soils, do not apply more than two weeks prior to planting. If an unsatisfactory length of weed control results from adverse environmental conditions following early treatment, a follow-up application of an appropriately labeled herbicide may be used. If the follow-up treatment includes atrazine, do not exceed the labeled rate for corn indicated in Table 1.

If weeds are present at the time of treatment, apply in tank mix combination with a contact herbicide (For example Gramoxone Extra or Roundup). Observe directions for use, precautions and restrictions on the label of the contact herbicide.

Note: To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated (broadleaf and Grass Control): Broadcast in the spring after plowing at the rate indicated in Table 1. Application may be made before, during or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation of LIQUID ATRAZINE 4L. Best results have been obtained when LIQUID ATRAZINE 4L is applied within two weeks prior to planting.

Preemergence or At-Planting (Broadleaf and Grass Control): Apply during or shortly after planting prior to weed emergence at the rate indicated in Table 1.

8/29/94

Postemergence (Broadleaf and Grass Control): Apply before weeds exceed 1.5 inches in height and before corn reaches 12 inches in height. For postemergence applications, if there has been no previous soil application to that crop, the maximum rate is 2 pounds a.i./A. If there has been a previous soil application to that crop, do not exceed total of 2.5 pounds a.i./A per calendar year. Apply at the rate indicated in Table 1.

TABLE 1 - For Control of Broadleaves and Grasses in Corn*

FOR ALL SOIL APPLICATIONS PRIOR TO CROP EMERGENCE

On Highly Erodible Soils (as defined by SCS)

If conservation tillage is practiced, resulting in at least 30% of the soil covered with plant residues at planting, apply a maximum of 2 pounds a.i./A (4 pts./A) as a broadcast spray.

If less than 30% of the soil is covered by plant residue at planting, a maximum of 1.6 pounds a.i./A (3.2 pts./A) may be applied.

On Soils that are Not Highly Erodible

Apply 4 pts./A as a broadcast spray.

FOR POSTEMERGENCE APPLICATION

If no atrazine was applied prior to corn emergence, apply a maximum of 4 pts./A broadcast.

If a postemergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.5 lbs. active ingredient (5 pts. of this product) per acre per calendar year.

* For preplant or pre-emergence applications (western Kansas, western Nebraska, eastern Colorado, eastern Wyoming, New Mexico, western Texas and the Panhandle of Oklahoma): on sands, loamy sands, sandy loams, mild to strong alkaline soils and all recently leveled soils, apply no more than 2.4 pints per acre for broadleaf weed control, either preplant surface, preplant surface incorporated, or preemergence. Many broadleaf weeds including pigweed, lambsquarters, nightshade, purslane and kochia will be controlled. On other soil types in the areas above, make applications at the rate shown in Table 1 for broadleaf and grass control.

Where there are state/local requirements regarding atrazine use (including lower maximum rates and /or higher set-backs) which are different from the label, the more restrictive/protective requirements must be followed. Certain states may have established rate limitations within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

Postemergence Broadleaf Weed Control with LIQUID ATRAZINE 4L Plus Emulsifiable Oil or Oil Concentrate in Water:

Broadcast 2.4 pints per acre for control of many broadleaf weeds including annual morning-glory, cocklebur, lambsquarters, mustard, pigweed, ragweed, smartweed and wild buckwheat. Application should be made before pigweed and lambsquarters reach 6 inches in height and before all other weeds reach 4 inches in height and before corn reaches 12 inches in height. Add emulsifiable oil at the rate of 1 gallon per acre for ground applications and one-half gallon per acre for aerial applications. A cultivation may be necessary if all weeds are not controlled or if regrowth of weeds occur

Postemergence Broadleaf and Grass Weed control with LIQUID ATRAZINE 4L Plus Emulsifiable Oil or Oil Concentrate in Water:

Broadcast 4 pints per acre after weed emergence, but before weeds reach 1.5 inches in height and before corn reaches 12 inches in height.

Add the following volume of one of the types oils indicated for aerial or ground application unless the oil label specifies otherwise

10 } 20
8

<u>Type Oil</u>	<u>Ground Application</u>	<u>Aerial Application</u>
Oil Concentrate (Crop or Petroleum-derived)	1 qt. / A	1/2-1 qt. / A
Petroleum-derived oil	1 gal. / A	2 qts. / A

Note: Crop-derived or petroleum-derived oil concentrates should contain at least 1%, but no more than 20% suitable emulsifier or surfactant blend.
 Petroleum-derived oil should contain at least 1% suitable emulsifier.

Precautions for Applications of LIQUID ATRAZINE 4L Plus Emulsifiable Oil in Water to Corn:

- (1) Do not use oils in sprays when treating inbred lines or any breeding stock as severe injury may occur.
- (2) Adding other insecticides, herbicides, liquid fertilizers or other materials is not recommended with Atrazine and emulsifiable oil in water because they may cause compatibility problems or crop injury.
- (3) Store and handle emulsifiable oil carefully. Oil contaminated with even a small amount of water may not emulsify properly when added to the tank.
- (4) Do not use oil in Atrazine sprays when crop is under stress from prolonged cold, wet weather, poor fertility, or other factors, or when crop is wet and succulent from recent rainfall, as crop injury may occur.
- (5) Do not exceed 2.5 lbs. active ingredient (or 5 pts. of this product) per acre per calendar year.
- (6) Postemergence application to corn must be made before corn exceeds 12 inches in height.

Tank-Mixtures with LIQUID ATRAZINE 4L on Corn:

This product may be tank mixed with these herbicides for control of certain broadleaf and grass weeds in corn:

- | | |
|----------------------------------|---------------------------|
| Alachlor | Gramoxone Extra |
| Alachlor + Gramoxone Extra | Princep |
| Alachlor + Roundup | Princep + Gramoxone Extra |
| Dual + Roundup (glyphosate) | Princep + Roundup |
| Dual (metolachlor) | Ramrod (propachlor) |
| Dual + Gramoxone Extra | Roundup |
| Dual + Princep | Sutan |
| Dual + Princep + Gramoxone Extra | |
| Dual + Princep + Roundup | |

Use tank mix directions appearing on the labels of the above herbicides when tank mixing with this product. Observe all precautions and limitations on labeling of products used in a particular tank mix.

When the labels of the above herbicides refer to atrazine 80W, use equivalent rates of Liquid Atrazine 4L. One pound of 80W equals 1.6 pints of 4L.

LIQUID ATRAZINE 4L Plus Gramoxone Extra - For control of existing vegetation and residual control where corn will be planted directly into a cover crop, established sod or in previous crop residues: Broadcast 4 pints LIQUID ATRAZINE 4L and 1-2 pints Gramoxone Extra per acre in 20-60 gallons of water per acre. Add 8 ounces of nonionic surfactant per 100 gallons of dilute spray. Add LIQUID ATRAZINE 4L to spray tank first and thoroughly mix with water. Add the Paraquat and surfactant last. Refer to Paraquat CL label for further directions, limitations and cautions.

LIQUID ATRAZINE 4L Plus Alachlor 4EC: Use a tank-mixture of LIQUID ATRAZINE 4L plus Alachlor 4EC for the control of many annual broadleaf and grass weeds in corn (field and silage corn only) including annual morning-glory, barnyard grass, black nightshade, Brachiaria, buttonweed (velvetleaf), carpetweed, cocklebur, crabgrass, fall panicum, Florida pusley, giant foxtail, green foxtail, yellow foxtail, goosegrass, lambsquarters, pigweed, purslane, mustard, common ragweed, smartweed and witchgrass.

Make applications at the rates indicated in Table 2. for preplant applications, apply within 7 days of planting and incorporate to a depth of 2-3 inches. For pre-emergence applications, apply to the soil surface after planting, but

11 9 20
7

before the crop or weeds emerge. For postemergence treatments, make applications until weeds reach the two-leaf stage, and the corn is no more than 3 inches tall.

Table 2. Tank-Mixtures of Liquid Atrazine 4L plus Alachlor 4EC on Corn (Field and Silage only)

Soil Texture	Broadcast Rate Per Acre			
	Less than 3% organic matter		More than 3% organic matter	
	Liquid Atrazine 4l (pts./A)	Alachlor 4EC (pts./A)	Liquid atrazine 4l (pts./A)	Alachlor 4EC (pts./A)
Light Sandy Soils	2	1.5	2	1.5
Silt Loam Soils	2-2.4	1.5-1.75	2-2.4	1.75-2
Heavy Clay Soils	2.4-3.2	2-2.5	2.4-3.2	2-2.5

NOTE: Apply in a minimum of 20 gallons of water per acre. Non-pressure fluid fertilizer may replace all or part of the water used as a carrier for application applied to the soil surface before crop emerges. Add the LIQUID ATRAZINE 4L to the spray tank first and thoroughly mix with water. Add the Alachlor last. At least 12 weeks must elapse following treatment with LIQUID ATRAZINE 4L Plus Alachlor 4EC before immature corn forage can be harvested or fed to cattle. Refer to the Alachlor 4EC label for further directions, limitations and cautions.

Princep 80W, Princep 4L or Princep Caliber®

In addition to the weeds listed under GENERAL DIRECTIONS for CORN, this combination also controls crabgrass, fall panicum, and carpetweed.

Broadcast tank mix before planting, at planting or after planting, but before crop and weeds emerge, at rates in table 2A. Use the 1:1 ratio for control of most weeds. Use the 1:2 ratio for expected heavy infestations of crabgrass and fall panicum. Cultivate shallowly if weeds develop.

Preplant Surface—Applied: Use on medium and fine textured soils with minimum-tillage or no-tillage systems only in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, WI and WY. Apply the recommended rate of Liquid Atrazine 4L and Princep shown in Table 2A up to 45 days prior to planting. Refer to the Liquid Atrazine 4L alone section for information if weeds should develop following the early treatment. On coarse-textured soils, do not apply more than 2 weeks prior to planting. Refer to the Liquid Atrazine 4L alone, preplant surface applied section under CORN section for additional details.

If weeds are present at the time of treatment, apply in a tank mix combination with a contact herbicide such as a paraquat product or Roundup. Observe directions for use, precautions, and restrictions on the label of the contact herbicide.

Note: To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

Preplant Incorporated: Apply to the soil and incorporate in the spring before, during, or after final seedbed preparation. Avoid deep incorporation. For best results, apply within 2 weeks prior to planting.

Preemergence: Apply during or shortly after planting, but before crop and weeds emerge. Refer to Corn Section of this label and to Princep 80w, Princep Caliber 90, or Princep 4L labels for further directions, limitations and precautions.

12920
7

Table 2A TANK MIXTURES WITH PRINCEP FOR WEED CONTROL IN CORN

Soil Texture	Broadcast Rate Per Acre			
	1:1 Ratio*		1:2 Ratio**	
	Liquid Atrazine 4L	Princep 80W ¹	Liquid Atrazine 4L	Princep 80W ¹
Sand, loamy sand, sandy loam	2 pts.	1.25 lbs	1.32 pts	1.67 lbs
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with low organic matter	2.4 pts.	1.5 pts	1.6 pts	2 lbs
Loam, silt loam, silt, clay loam, sandy clay loam, silty clay loam, sandy clay, or silty clay with medium to high organic matter, and clay (including dark prairie soils of the Corn Belt)	3 pts.	1.8 lbs	1.92 lbs	2.4 lbs

*For control of most weeds

** For control of expected heavy infestations of crabgrass and fall panicum.

¹ When using Princep Caliber 90 or Princep 4L, use equivalent rates. One (1) lb of Princep 80 W = 0.9 lb of Princep Caliber or 1.6 pts of Princep 4L.

Princep 80W, Princep 4L, or Princep Caliber 90 plus Roundup: Use as tank mixture for preemergence and postemergence control of certain broadleaf and grass weeds where corn will be planted directly into a cover crop, established sod, or in previous crop residues. Refer to Roundup label for all directions, weeds controlled, precautions and limitations.

Princep 80W, Princep 4L, or Princep Caliber 90 plus Gramoxone Extra: Use as tank mixture with Princep and Gramoxone Extra to kill existing vegetation and for residual weed control where corn will be planted directly into a cover crop, established sod, or previous crop residues. Add this product and princep to water in spray tank, agitating until thoroughly mixed. Then add Gramoxone Extra or Paraquat CL and a nonionic surfactant, such as X-77. Continue agitation during application. Broadcast 2-4 pts of this product plus 1.25-2.5 lbs of Princep 80W (or 2-4 pts of Princep 4L, or 1.1 - 2.2 lbs of Princep Caliber 90) plus the recommended labeled rate of Gramoxone Extra for this tank mixture, in 20-60 gallons of water per sprayed acre. Apply before during or after planting, but before the corn emerges. Add 0.5 pts of a nonionic surfactant such as X-77 per 100 gallons of spray mixture. Use the higher rate of Gramoxone Extra specified on the label if existing vegetation is 4-6 inches tall. This mixture will not control weeds taller than 6 inches.

Refer to further limitations and precautions on the labels for this product, Princep and Gramoxone Extra or Paraquat CL.

Precautions for all Applications to Corn:

- (1) The maximum application rate for corn is 2.5 lbs. active ingredient per acre per calendar year. Therefore, do not apply more than 5 pints of LIQUID ATRAZINE 4L on corn in any one year.
- (2) Following Harvest of a treated crop, plow (moldboard or disk-plow) and thoroughly till the soil in the fall or spring to minimize possible injury to rotational spring-seeded crops, regardless of the rate used.
- (3) For best control of velvetleaf and cocklebur, the application rate cannot be less than 2 pounds per acre of active ingredient, either alone or in tank mix combinations.
- (4) Do not graze treated area or feed-treated forage to livestock for 21 days following application of this product.

For postemergence applications plus emulsifiable oil, see additional precautions under that section of the label.

Sorghum and Sorghum-Sudan Hybrids (Grain and Forage Types): LIQUID ATRAZINE 4L may be applied either before planting or after planting as indicated below. For soil applications prior to crop emergence (i.e., early preplant, preplant incorporated, preplant surface, at planting or pre-emergence) see table 1.

Preplant Incorporated (Broadleaf and Grass Control): Broadcast in the spring after plowing at the rate indicated in Table 3. Application may be made before, during or after final seedbed preparation. If soil is tilled or worked after application, avoid deep incorporation of LIQUID ATRAZINE 4L. For best results, apply within two weeks prior to planting.

Preplant Surface Applied (Broadleaf and Grass Control): Use on medium and fine textured soils with minimum tillage systems only in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, WI and WY. Apply the recommended rate of LIQUID ATRAZINE 4L shown in Table 3 up to 45 days prior to planting. If an unsatisfactory length of weed control results from adverse environmental conditions following early treatment, a follow-up application of appropriately labeled herbicide may be used. If the follow-up treatment includes atrazine, do not exceed the labeled rate for corn indicated in Table 1. Under dry conditions, irrigation after application is recommended to move Liquid Atrazine 4L into the soil.

If weeds are present at time of treatment, apply in a tank mix combination with a contact herbicide such as Gramoxone Extra or Paraquat CL, or Roundup. Observe directions for use, precautions and restrictions on the label of the contact herbicide.

Note: To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting or weed control will be diminished.

Preemergence or At-Planting (Broadleaf and Grass Control): Apply during or shortly after planting, but prior to weed or crop emergence at the rate indicated in Table 3.

Post-emergence (Broadleaf and grass control): Apply at rate shown in Table 3. before weeds exceed 1.5 inches in height, and before sorghum is taller than 12 inches.

TABLE 3 - Broadleaf and Grass Weed Control in Sorghum ^{(1),(2)}

<p>FOR ALL SOIL APPLICATIONS PRIOR TO CROP EMERGENCE</p> <p><u>On Highly Erodible Soils</u> (as defined by SCS)</p> <p>If conservation tillage is practiced, leaving at least 30% of the soil covered with plant residues at planting, apply a maximum 2 pounds a.i./A (4 pts./A) as a broadcast spray.</p> <p>If the soil coverage with plant residue is less that 30% at planting, a maximum of 1.6 pounds a.i./A (3.2 pts./A) may be applied.</p> <p><u>On Soils Not Highly Erodible</u></p> <p>Apply 4 pts./A as a broadcast spray.</p> <p>FOR POSTEMERGENCE APPLICATION</p> <p>If no atrazine was applied prior to sorghum emergence, apply a maximum of 4 pts./A broadcast. If a postemergence treatment is required following an earlier herbicide application, the total atrazine applied may not exceed 2.5 lbs. active ingredient (5 pts. of this product) per acre per calendar year.</p>
--

⁽¹⁾ LIQUID ATRAZINE 4L should not be applied preplant surface or preplant incorporated to sorghum grown in AL, AR, FL, GA, LA, MS, NC, NM, OK, SD, TN, or TX. Do not apply pre-emergence to sorghum grown in NM, OK, or TX, except in northeast OK and the TX Gulf Coast and Blacklands areas.

⁽²⁾ Where there are state/local requirements regarding atrazine use (including lower maximum rates and /or higher set-backs) which are different from the label, the more restrictive/protective requirements apply. Certain states may have established rate limitations within specific geographical areas. consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.

In case of planting failure, sorghum or corn may be replanted into soil previously treated with LIQUID ATRAZINE 4L. Do not make a second broadcast application or injury may occur. If LIQUID ATRAZINE 4L is applied in a band and sorghum is replanted in the untreated row middles, LIQUID ATRAZINE 4L can be applied in a band to the second planting provided the maximum application rate of 2.5 ai/A per calendar year is not exceeded.

Preemergence Broadleaf Weed Control in Furrow Irrigated, Bedded Sorghum (AZ and CA only): For preemergence control of many broadleaf weeds, broadcast 1.5 - 2 pts /A after bed preparation, during or after planting, but before sorghum and weeds emerge and before the first furrow irrigation. Follow with several regular irrigations, making sure to thoroughly wet all soil.

Precautions for Preemergence Application to Furrow Irrigated Bedded Sorghum in AZ and CA: To avoid possible sorghum injury, do not use on sand or loamy sand soil or on sorghum planted in furrows. Application to sorghum growing on alkali soils or where cuts, fills, or erosion have exposed calcareous or alkali subsoils may cause crop injury. In case of crop failure, do not replant sorghum for 8 months following application. Corn may be planted immediately.

Postemergence Broadleaf Weed control with LIQUID ATRAZINE 4L Plus Emulsifiable Oil or Oil Concentrate in Water: Broadcast 2.4 pts./A for control of broadleaf weeds including annual morning-glory, cocklebur, lambsquarters, mustard, pigweed, ragweed, smartweed and wild buckwheat. Application should be made before pigweed and lambsquarters reach 6 inches in height and before all other weeds reach 4 inches in height. In CO, western KS, NM, OK, TX, and other desert regions of AZ and CA, apply when sorghum is 6-12 inches in height, but before it reaches the boot stage. In all other areas, apply after sorghum reaches 3-leaf stage and before it exceeds 12 inches in height. Add emulsifiable oil at the rate of 1 gallon of emulsifiable oil / A for ground applications and 0.5 gal / A for aerial applications. Add 1 qt / A of oil concentrate for ground application. A cultivation may be necessary if all weeds are not controlled or if regrowth of weeds occurs.

For a more complete list of the weeds controlled, see "Broadleaf and Grass Weeds Controlled with Atrazine Alone" Under "GENERAL INFORMATION"

Precautions for Applications of LIQUID ATRAZINE 4L Plus Emulsifiable Oil in Water to Sorghum: See precautions under "Atrazine Plus Emulsifiable Oil or Oil Concentrate in Water Applications to Corn".

Postemergence Broadleaf Weed Control with LIQUID ATRAZINE 4L Plus Surfactant (CO, western KS, NM, OK, TX, and desert regions of AZ and CA only): Broadcast 2.4 pts. / A LIQUID ATRAZINE 4L plus .75 to 1.5 pts. of surfactant after sorghum reaches 6 inches in height, but before weeds reach 1.5 inches in height. Apply only on sandy loam and finer textured soils.

Precautions for All Applications to Sorghum

- (1) Heavy rains immediately following application tend to result in excessive concentrations of herbicide in seed furrow, resulting in possible crop injury. Applications to furrow-planted sorghum should not be made until furrows are leveled (plow-in). Deep planter marks or seed furrows should also be leveled before application.
- (2) Application made to sorghum growing under stress caused by minor element deficiency or to sorghum growing on highly calcareous soils may result in crop injury.
- (3) Following harvest of a treated crop, plow (moldboard or disk-plow) and thoroughly till the soil in the fall or spring to minimize possible injury to rotational spring-seeded crops, regardless of rate used.
- (4) Crop injury may occur when this herbicide is either applied preplant incorporated or pre-emergence in conjunction with an at-planting systemic insecticide.
- (5) The maximum application rate for sorghum is 2.5 lbs. active ingredient (5 pts. of this product) per acre per calendar year.
- (6) For all soil applications prior to crop emergence (except for preemergence use on bedded sorghum in AZ and CA), do not apply to coarse-textured soils, i.e., sand, loamy sand, sandy loam, or to medium- and fine-textured soils having less than 1% organic matter, or injury may occur.
- (7) For postemergence applications, do not apply to sand or loamy sand, or injury may occur.

Note: Do not graze or feed forage from treated areas for 21 days following application.

8/29/94

Rotational Crop Suggestions: See "rotational crop" recommendation under corn.

Tank Mixtures For Grain Sorghum

Dual 8E: Use as a tank mixture with Dual 8E for control of those weeds listed on the Dual 8E label, as well as on this label. Use this tank mixture only on sorghum seed treated with Concep. Refer to the Dual 8E label for all directions, precautions and limitations.

WINTER WEED CONTROL IN TEXAS

For postemergence control of winter weed only, such as henbit, seedling dock, and annual thistle on fall bedded land in the Gulf Coast and Blacklands of Texas. Apply 0.8 - 1 qt / A postemergence to the weeds in November or December to land that will be planted to corn, grain sorghum or forage sorghum the following spring. For best results, add a suitable surfactant, such as X-77, at the rate of 0.5% of the spray volume, or an emulsifiable oil at the rate of 1.0% of the spray volume.

Normal weed control programs may be used in the following corn, grain sorghum, or forage sorghum crop.

NOTE: Do not plant any crops except corn, grain sorghum or forage sorghum the spring following this treatment, or illegal residues may result.

CHEMICAL FALLOW - LIQUID ATRAZINE 4L

Wheat-Fallow-Sorghum: To control annual broadleaf and grass weeds following wheat harvest and in the following sorghum crop, when grown under minimum tillage, broadcast 6 pts. / A to wheat stubble immediately following wheat harvest. If weeds are present at application, remove them with a sweep plow or other suitable implement after application, or used an approved contact herbicide before or after application of LIQUID ATRAZINE 4L. Plant sorghum into wheat stubble the following spring with minimum soil disturbance. Use a surface planter or a planter leaving a shallow furrow. If weeds are present at sorghum planting, remove them with a sweep plow or other suitable implement before planting.

For the list of weeds controlled, see the list under "GENERAL INFORMATION"

Precautions for Weed Control in Chemical Fallow: (1) Use only on silt loam or finer textured soil, or crop injury may result. (2) Wheat -Fallow-Sorghum cropping sequence must be followed. (3) Do not apply following sorghum harvest.

NOTE: Do not graze or feed forage from the treated area to livestock. To avoid illegal residues and crop injury, do not plant any crop other than those on this label within 18 months following treatment.

Wheat-Fallow-Corn (CO, KS, ND, NE, SD and WY): This product controls cheat grass (downy brome, chess), kochia, mustards, pigweed, Russian thistle, wild lettuce, wild sunflower, and volunteer wheat during the period after wheat harvest. Weed control may extend into the following corn crop grown under minimum tillage.

On soils in ND and SD with a pH greater than 7.5, do not exceed 1.5 lb a.i. / A of Liquid Atrazine 4L. For soils with a pH less than 7.5 in ND and SD, apply 1-2 lb a.i. / A. Use the higher rate on finer textured soils and where heavy weed infestations are expected. Use the lower rate on coarse textured soils and where light weed infestations are expected. In the event grasses are present in the following spring, use a grass herbicide registered for use on corn.

Follow directions for use, notes and precautions in the Wheat-Fallow-Sorghum section above, substituting corn where references to sorghum are made.

Wheat-Fallow-Wheat (CO, KS, ND, NE, SD and WY): For preemergence control of cheatgrass (downy brome, chess), common lambsquarters, field pennycress, kochia, mustard, Russian thistle, wild lettuce, and suppression of volunteer wheat during fallow period of a wheat-fallow-wheat rotation, broadcast 1-2 pts. / A on all soils except those listed under Precautions. For control of pigweed and wild sunflower, use only the higher rate. Apply to stubble ground. Treat only once during the same fallow period.

TANK MIXTURES FOR CHEMICAL FALLOW

Wheat-Fallow-Sorghum or Wheat-Fallow-Corn (KS, NE)

Gramoxone Extra: If weeds are present at application, a tank mix with Gramoxone Extra may be used. Broadcast 6 pints of Liquid Atrazine 4L plus a suitable amount of Gramoxone Extra in 20-60 gallons of water per acre by ground equipment. Add 0.5-1 pint of a nonionic surfactant, such as X-77, per 100 gallons of spray mixture. Add this product to the spray tank first and thoroughly mix with water. Then add Gramoxone Extra followed by surfactant. Use the higher rate of Gramoxone Extra specified on label if weeds are 4-6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during the same fallow period. Refer to label for Gramoxone Extra for further directions, precautions and limitations.

Wheat-Fallow-Wheat (CO, KS, ND, NE, SD, and WY)

Gramoxone Extra: If weeds are present at application, a tank mix with Gramoxone Extra may be used. Broadcast 1-2 pints of Liquid Atrazine 4L plus a suitable amount of Gramoxone Extra in 20-60 gallons of water per acre by ground equipment. Add 0.5-1 pint of a nonionic surfactant, such as X-77, per 100 gallons of spray mixture. Add this product to the spray tank first and thoroughly mix with water. Then add Gramoxone Extra followed by surfactant. Use the higher rate of Gramoxone Extra specified on label if weeds are 4-6 inches tall. This mixture will not control weeds taller than 6 inches. Apply to stubble ground. Treat only once during the same fallow period. Refer to label for Gramoxone Extra for further directions, precautions and limitations.

If weeds are present at application of this product alone, use an approved contact herbicide either before or after treatment, or tillage after treatment.

Use tillage to control weeds which escape during fallow period. Till before planting. For this product applied alone or in tank mixture Gramoxone Extra, plant at least 2 inches deep and 12 months or more after application.

Precautions: To avoid crop injury, (1) do not use on sand soil. (2) Do not treat eroded hillsides, caliche and rocky outcroppings, or exposed calcareous subsoil. (3) Do not treat soils of the Rosebud and Canyon series in western Nebraska and adjoining counties in CO and WY. (4) Do not treat soils with calcareous surface layers. (5) Avoid spray overlap.

Note: Do not graze treated areas within 6 months after application.

Aerial Application: In order to assure that spray will be controllable within the target area when used according to label directions, make applications at a maximum height of 10 ft, using low drift nozzles at a maximum pressure of 40 psi., and restrict application to periods when wind speed does not exceed 10 mph. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply Liquid Atrazine 4L by aircraft at a minimum upwind distance of 400 ft from sensitive plants.

CONSERVATION RESERVE PROGRAM (CRP) (NE, OK, OR AND TX)

For control or suppression of the following weeds: annual ragweeds, barnyardgrass, black nightshade, cheat, cocklebur, downy brome, fall panicum, field pennycress, giant foxtail, yellow foxtail, Japanese brome, Kentucky bluegrass, kochia, lambsquarters, little barley, maretail, pigweed, prickly lettuce, smooth brome, and sunflower, refer to directions, notes, and precautions below.

Pure stands of newly seeded big bluestem, switchgrass and eastern gamagrass.

Use only on loam, silt loam, silty clay loam, clay loam, and silty clay soils with at least 1% organic matter.

Establishment: Broadcast 2 qt./A of Liquid Atrazine 4L, preplant incorporated or preemergence, at time of seeding and prior to emergence of weeds. Prepare a good firm seedbed. Plant 1/2 inch deep with a grassland drill (preferred method) or a conventional drill. If a conventional drill is used on prepared seedbeds, remove all tension from the disk openers. For best results, cutback or roll after planting.