



# RABON® ORAL LARVICIDE BLOCK

To prevent the development of horn flies, face flies, house flies and stable flies in the manure of treated cattle.

## ACTIVE INGREDIENT

2-Chloro-1(2, 4,5-Trichlorophenyl)  
Vinyl Dimethyl Phosphate ..... 0.463%  
Inert Ingredients\* ..... 99.537%

\*REFERS ONLY TO INGREDIENTS WHICH ARE NOT INSECTICIDAL

## GUARANTEED ANALYSIS

Calcium (Ca), not more than ..... 13.0%  
Calcium (Ca), not less than ..... 11.0%  
Phosphorus (P), not less than ..... 4.0%  
Sodium (NaCl), not more than ..... 21.0%  
Salt (NaCl), not less than ..... 18.0%  
Iodine (I), not less than ..... 0.0003%  
Vitamin A, USP Units Per Pound, Min ..... 60,000

## INGREDIENTS

Dicalcium Phosphate, Monocalcium Phosphate, Ground Limestone, Salt, Processed Grain By-Products, Cane Molasses, Lignin Sulfonate, Vitamin A Supplement (Stability Improved), D-Activated Animal Sterol (Source of Vitamin D<sub>3</sub>), Vitamin E Supplement, Magnesium Oxide, Potassium Chloride, Calcium Sulfate, Manganous Oxide, Iron Carbonate, Iron Oxide, Zinc Oxide, Copper Oxide, Cobalt Carbonate, Potassium Iodide

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS

Keep out of reach of children. Harmful if swallowed. Avoid contact with skin and eyes. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating or smoking. If in eyes, wash with plenty of water for 15 minutes. If irritation persists, see a physician.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

EPA Reg. No. 01990-387  
EPA Est. No. 01990-KS-01

**NET WEIGHT 50 LBS.**

Rabon is a registered trademark of Shell Oil Company.

**MANUFACTURED BY  
FARMLAND INDUSTRIES, INC.  
GENERAL OFFICES: KANSAS CITY, MO. 64116  
81-7198 (Series 08KS)**

## DIRECTIONS FOR USE GENERAL CLASSIFICATIONS

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

1. To be fed daily at the rate of .07 grams of larvicide in 0.5 ounce of block per 100 pounds of body weight. Example: 5 ozs. (approximately 1/4 pound) of block for a 1000-pound animal daily.
2. Provide at least one block per five head of cattle. (So timid and smaller animals can get their share.)
3. Place the blocks where animals congregate (loafing, grazing, feeding and watering area).
4. Feed blocks continuously. Add one additional block when EACH block has been ONE-HALF consumed.
5. Do not feed salt or mineral supplements of any kind containing salt.
6. For initial control of adult flies, spray your cattle to reduce the existing population. The normal daily intake of CO-OP Rabon® Oral Larvicide Block will then prevent the development of new fly larvae in the manure of treated cattle. Because face flies migrate, work with your neighbors to more effectively control the adult face fly population. This product is not effective against existing adult flies. Supplemental fly control measures may be needed in and around cattle lots and buildings to control adult house flies and stable flies which can breed not only in cattle manure but in other decaying vegetable matter or silage on the premises. In order to achieve optimum fly control, CO-OP Rabon® Oral Larvicide Block should be used in conjunction with other good management and sanitation practices.

## STORAGE AND DISPOSAL

Do not contaminate water or feed by storage or disposal. Open dumping is prohibited. Pesticide that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

**ACCEPTED**

**MAR 20 1981**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 1990-387

20