


GOOD-WAY

Rotenone Dust

1.50% Rotenone

CONTROLLING Cattle
Grubs, Cattle Lice, Sheep
AND MANY OTHER
INSECTS
TREATS A POUNDS

ACTIVE INGREDIENTS:
Rotenone 1.50%
Other
Cube Resins 4.77%
INERT INGREDIENTS 93.73%
100.00%



ACCEPTED
June 1963
UNDER THE
FUNGICIDE ACT
FOR ECONOMIC
ED UNDER NO

DIRECTIONS:

CATTLE GRUBS: Apply dust liberally "around grub welts," using a stiff brush, and scrub powder into grub holes.

Repeat every thirty days, during Winter and Early Spring months if grub bumps are present!

CATTLE LICE: Apply dust liberally over back from tailhead of animal to nostrils, also other parts of body that might be infested, such as udder flanks and inside of legs. Dust should be well brushed into the skin.

A second treatment should be made in 12 to 14 days and if necessary, follow with a third treatment within 3 weeks after the first treatment.

SHEEP TICKS: Dust sheep immediately after shearing, using hand dust gun or powder duster. One pound of Rotenone Dust will treat 8 animals.

Repeat within 24 days if necessary. Lambs should be dusted as soon as ticks appear.

IN THE GARDEN: Use Rotenone Dust for dusting garden plants, especially good for controlling pea aphids, imported cabbage worms, cabbage loopers, cucumber beetles, and spittlebugs on strawberries.

Thoroughly dust plants when insects appear. Repeat as necessary.

Distributed by
GOOD-WAY INSECTICIDES, INC.
WHEELING, ILLINOIS

WARRANTY

NOTICE: Good-Way Insecticide, Inc. makes no warranty, express or implied, concerning this material except that it conforms to the chemical description on the label.

Neither Good-Way Insecticide, Inc. nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting from the handling, storage or use of this material. The buyer assumes all risk and liability therefrom and accepts and uses this material on these conditions.

2006-36

06/06/63

1/1