

WARNING! HAZARDOUS IF SWALLOWED, INHALED, OR ABSORBED THROUGH SKIN

Wash thoroughly with soap and water after handling and before eating and smoking. Wear clean clothing. In case of spillage on person or clothing, immediately remove clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Keep away from heat and open flame. ATROPINE IS ANTIDOTAL.

PRECAUTIONS IN USING

Keep out of the reach of children and domestic animals. Do not contaminate feed or foodstuffs. During commercial or prolonged exposure in spray-mixing and loading operations, wear clean rubber gloves and a mask or respirator of a type found suitable by the U. S. Department of Agriculture for protection against Ciodrin Insecticide. Use this material only for recommended purposes and at recommended dosages.

DECONTAMINATE AND DESTROY USED CONTAINERS

Do not re-use for any purpose. Decontaminate containers by rinsing thoroughly with water and alkaline detergent. Destroy containers by breaking and burying fragments in isolated area. Dispose of rinsings in a way as not to constitute a hazard, or contaminate water supplies.

Distributed by
GOOD-WAY INSECTICIDE, INC.
Wheeling, Illinois
U. S. D. A. Reg. No. 2006-37

GOOD-WAY

CIODRIN*

DAIRY LIVESTOCK SPRAY BACK RUBBER CONCENTRATE

Contains 1.1 lb. Active Ingredient
Per Gallon.

ACTIVE INGREDIENTS: BY WEIGHT	
Dimethyl phosphate of Alpha-methylbenzyl 3-hydroxy-cis-crotonate	14.4%
Petroleum hydrocarbons	73.0%
INERT INGREDIENTS: 12.6%	
Total	100.0%

*Ciodrin Insecticide
*Trademark Shell Chemical Company
USDA Reg. No. 2006-37

INSECTICIDE EMULSIBLE CONCENTRATE

Approved for use on Dairy and Beef Cattle, Horses, Hogs, Sheep, Goats. Controls: Stable flies, Horn flies, House flies, Lice, and Lone Star Ticks.



WARNING
Keep Out of Reach of Children and Domestic Animals

GENERAL DIRECTIONS

CIODRIN Insecticide Emulsible Concentrate is suitable for use in conventional hydraulic sprayers or low pressure knapsack sprayers. Prior to use, mix the required amount in sufficient water and apply in accordance with State and Federal Recommendations. Do not apply regularly to calves under 6 months of age. Brahmen Cattle should not be treated as they may show hypersensitivity to organic phosphates.

DIRECTIONS FOR LIVESTOCK PEST CONTROL

For control of stable flies, horn flies, house flies, lice and lone star ticks on dairy cattle, beef cattle, horses, sheep, goats and hogs—Mix 2 qts. of this product per 6 gals. of water (1% conc.). Spray to thoroughly cover all parts of the animal, including the legs, using 1-2 pts. of spray per large animal and proportionately less for smaller animals. Repeat application as necessary to maintain control but not more often than 7 days.

For pen or corral operations involving spray applications to livestock with high pressure equipment, do not exceed a 0.5% conc.—Mix 2 qts. of this product per 12 gals. of water (0.5% conc.). Spray to thoroughly cover the animals body and legs (avoid excessive spraying of the head) using 1-2 qts. of spray per large animal and proportionately less for smaller animals.

Improved face fly control may be achieved by adding sugar to the spray solution at the rate of 1 lb. per 6-10 gals. of spray.

Low Volume Application for Control of Face Flies—Mix 4 Tbsps. (2 fl. oz.) of this product per 1-1/2 pts. of water (1% conc.). Using a calibrated sprayer, apply 5 tsps. of spray to the face and total of 5-10 tsps. to the back and sides of the animals 3 times per week. Improved control may be achieved by the addition of 1 tsp. of sugar per pt. of spray solution.

For Control of House Flies, Face Flies, and Stable Flies in and Around Livestock Buildings—Mix 2 qts. of this product per 6 gals. of water (1% conc.). Spray premises such as inside and outside barn walls (including dairy barns), eaves, fences and other resting or breeding places at least twice monthly using approximately 1 gal. of finished spray per 1000 sq. ft. Improved house fly and face fly control may be achieved by adding sugar to the spray solution at the rate of 1 lb. per 6-10 gals. of spray.

For Control of Stable Flies, Horn Flies, House Flies, and Face Flies on Dairy and Beef Cattle, and Horses—Mix 8 tbsps. (4 fl. oz.) of this product per 1 1/2 pts. of water (2% conc.). Spray to thoroughly cover all parts of the animal, including the legs, using 1-2 fl. oz. of spray per animal. Apply daily with conventional hand or power sprayers. Improved control may be achieved by the addition of 1 tsp. of sugar per pt. of spray solution.

For Control of Lice and Winter Ticks on Dairy Cattle, Beef Cattle, Horses, Sheep Goats and Hogs—Mix 1-2-1/2 pts. of this product per 16 gals. of water (0.1%-0.25% conc.). Spray animals thoroughly using up to 2 gals. of spray for the 0.1% conc. and up to 1 gal. of spray for the 0.25% conc. Apply a second application 14 days later. Repeat applications as necessary to maintain control.

Backrubber Application: For use in backrubbers or fare rubbers for the control of horn flies and face flies, mix 1 qt. of this product per 4 gals. of back rubber base oil (1% conc.). This may be diesel fuel or fuel oil in accordance with local recommendations.

APPROVED
 Apr. 29, 1970
 2006-37